

Malayan


Wason
SH
117
M2
M46


ASIA

Fishes

by

C. N. Maxwell.





Cornell University Library

The original of this book is in
the Cornell University Library.

There are no known copyright restrictions in
the United States on the use of the text.

Cornell University Library
SH 117.M2M46

Malayan fishes /



3 1924 006 251 569

ech


Malayan Fishes

by

C. N. MAXWELL

Director of Supplies, S. S. and F. M. S.

Singapore
Printed at the Methodist Publishing House
1921.



Contents.

	Page
1. Preface	3
2. Introduction	5
3. Part I. Descriptions of Species	15
4. Part II. Malay Vocabulary of Fishes	54
5. Part III. Systematic Classification of the Fishes described	86
6. Authors consulted	102
7. Index to English names	105
8. List of Plates	107
9. Plates.	

Preface.

Literature dealing with our local Fish and Fisheries is wanting.

The members of the Profiteering Commission (1920) who were "impressed and much handicapped by the absence of any recorded information relating to fish and fisheries," made certain recommendations for the future control and organisation of the fishing industry and these recommendations have received the approval of the Government.

With the best will in the world, the task of Legislators and Fishery Officers who have no data or records to guide them, and are therefore unable to discuss our local fish and fisheries except in vague terms, would be as fruitless in the future as it has been in the past.

Allowing, therefore, that recorded information in the form of a hand-book on Malayan Fishes is wanted and wanted at once, the difficulty arises that there is no one qualified or likely to be qualified for some years to write such a book.

The ichthyologists are not linguists and the linguists are not ichthyologists.

This being the position, the writer has the temerity to offer this small work, which he hopes will be of some temporary service until, in due course, the importance of the Malayan Fisheries has been established and Fishery bulletins written by specialists are produced.

The inclusion in this volume of several hundred Malay names of fishes, many published for the first time, should lighten the labours of scientists and help the Fishery Officers.

No fishes have been included which have not been definitely recorded as inhabiting the seas, estuaries and fresh water of the Malay Peninsula.

The size of the work would have been trebled if fishes of Borneo, Java, Sumatra and the Malay Archipelago generally, together with Siam and Burma, had been admitted. It is safe to prophesy that most of the fishes of those countries inhabit our waters and will be recorded later on.

This work may be taken, therefore, as dealing, very inadequately, with one-fourth of our fishes only and probably not one-half of the local Malay names have been mentioned. The writer knows very little about fresh water fishes.

The material in this book has been put together hastily during a period of five months in the intervals of considerable pressure of other work.

The plates have been beautifully prepared by Mr. Black of the Survey Department, Kuala Lumpur, but it is to be regretted that many of the fishes have been badly displayed and badly photographed.

The writer was unable to find time to be present at the Clyde Terrace Market, Singapore, where most of the photographs were taken, the fish being borrowed for a minute or two from the stallholders, and in consequence, the specific identification of every fish from a poor photograph has been impossible, though the writer feels confident that the families and genera have been correctly given.

The writer's thanks are due to Messrs. Stead and Roughley. But for their works on Australian Fishes, from which quotations have been freely made, this work could not have been written.

To the Directors and Staff of the F. M. S. Museums and the Raffles Museum, Singapore, who have granted me facilities for consulting the reference libraries and permission to examine and photograph specimens in the Museum collections, I desire to express my indebtedness.

C. N. MAXWELL,
Director of Supplies.

Singapore, 16th June, 1921.



Malayan Fishes

BY

C. N. MAXWELL

INTRODUCTION.

"Fish is not a luxury, but an absolute necessary of life, with a rice-eating population."

"It is obvious that in order to secure an adequate and plentiful supply of fish, especially to large cities like Calcutta.....we must go further out—into the deep sea—which, after all, is the largest repository of piscine wealth.....facts and figures relating to the sea-fisheries of Great Britain, the United States and Canada.....ought to open our eyes to the great possibilities which lie before us."

"In Bengal, Government will have to do a great deal more; it will have to *create* and *build up* the sea-fishing industry, with the object of handing it, let us hope at no distant date, to private enterprise.

"It will also be necessary to show the best way of working the estuarine fisheries by improved methods of capture and of bringing the catches expeditiously to market in a sound state."

Sir K. Gupta, K. C. S. I. Report on Fisheries of Bengal and into Fishery matters in Europe and America, 1908.

"I appeal to the whole population of these Islands, a maritime people who owe everything to the sea. I urge them to become better informed in regard to our national sea-fisheries and take a more enlightened interest in the basal principles that underlie a rational regulation and exploitation of these important industries. National efficiency depends to a very great extent upon the degree in which scientific results and methods are appreciated by the people and scientific investigation is promoted by the Government and other administrative authorities. The principles and discoveries of science apply to aquiculture no less than to agriculture. To increase the harvest of the sea the fisheries must be continuously investigated"

W. A. Herdman, C.B.E., D. Sc., F.R.S., etc. Annual address of the President of the British Association 1920.

"In no other section of our food supply.....could the application of capital to a comparatively small amount mean so considerable a development.....Both as regards railway and cold storage facilities the fish trade is in its infancy.....Transportation—cheap and rapid, must be provided by the State—fish trains should have precedence—and rates should be very low, even to the extent of entailing considerable loss."

The Earl of Dunraven. Paper read before the Royal Statistical Society, March 20, 1917.

Fish are curious creatures and we have still a great deal to learn about their habits. Some like the Salmon and the Shad (**Ikan t rubok**) live in the sea and spawn in the rivers. Such fish are termed anadromous and the term is also applied to fish which make a migration from the deep sea coastwards for the purpose of spawning.

Others, like some Eels, live in the rivers and spawn in the sea. The common Eel of Europe (*Anguilla vulgaris*) spawns far out in the ocean, after which both males and females die, never returning to fresh-water a second time. Fishes which live in the rivers and spawn in the sea are termed catadromous.

Some fishes do not lay eggs but bring forth their young alive. Examples of viviparous fishes occur in the Shark and Ray families and also in the BLENNIDAE, CYPRINODONTIDAE and SCORPAENIDAE. Instances of functional hermaphroditism occur, and some of the SERRANIDAE (Sea-Perches) are invariably hermaphrodite and self-fertilising.

A Sea-Bream, *Chrysophrys auratus*, is an example of successive hermaphroditism, the male and female sex-cells ripening alternately. As an occasional variation hermaphroditism has been recorded in such well known fishes as the Cod, the Mackerel and the Herring.¹

The eggs of fishes may be divided into two kinds; the large (*demersal* ova) which are heavy and sink; and the small (*pelagic* ova) which are buoyant and float at or below the surface according to their density. The buoyancy of the pelagic egg depends, however, on the density of the sea and the pelagic egg becomes demersal, in position, in brackish water and in fresh water.

Demersal eggs may be either viscid and adhesive or smooth and non-adhesive.

Pelagic eggs are distinguished by their lightness, buoyancy, small size and remarkable transparency. They are always non-adhesive and free and they invariably belong to Marine Fishes. As a general rule it may be said that fresh water fish produce demersal ova and marine fish pelagic ova.

When we realise that the eggs of most Marine fishes float, it is obviously futile to speak of guarding the "spawning grounds" on our coasts. It is necessary to mention this because at one time it was thought that spawning took place on shallow banks or even close in shore but this is now known to be incorrect, except in the case of the true Herring which lays demersal eggs in comparatively shallow water, and a few less important species.

Amongst our important Marine food fishes which are known to produce pelagic eggs are members of the Herring, Mackerel, Horse-Mackerel, Sea-Perch, Mullet and Flatfish families, in fact, all our best fish.

¹ Camb: Nat: Hist: 1904.

Fishes known to produce demersal eggs on our coasts are the Gar-Pike (**Todak**) and the Flying-fish (**Bélalang**) and their eggs have viscid threads by which they become attached or entangled with foreign objects or eggs of the same species. The eggs of the **Todak** may be seen entangled in fishing stakes (*kelong*) in masses, which look rather like cobwebs.

When the breeding season arrives fishes migrate to the localities most suitable for the deposition of their eggs. At this time our principal food fish which produce pelagic eggs proceed far out to sea against the prevailing monsoonal current. This is known as the contranantant spawning migration. After spawning, the eggs are brought back by the current towards the coast. This is the denatant drift.

Though the eggs of many species of fish hatch out fry which are miniature representations of the adult fish, the eggs of others hatch out larval forms, known as Leptocephali, which bear no resemblance to their parents. These Leptocephali are transparent, attenuated creatures, often ribbon-like in shape, with very small heads. They appear to be incapable of much effort and to be specially adapted for passive drift; in fact, the Leptocephalus stage appear to be a marvellous provision of Nature to enable the young of certain fish which spawn far out at sea to reach the shallows near the coasts in a state of suspended animation. We know that the Tarpin (*Megalops cyprinoides*) Malay **Bulan-bulan** and the Giant Herring (*Elops hawaiiensis*) Malay **Bandang**, pass through a Leptocephalus stage, and as no Malay fisherman whom I have questioned, has ever seen the **Parang-parang** (*Chirocentrus dorab*) until it was a few inches long, it may be because this fish passes through a larval metamorphosis also. It is only within recent years, that certain Leptocephali, long known to naturalists, have been identified as larval Eels.¹

For example, *Leptocephalus brevirostris* is now known to be the larva of the common Eel of Europe (*Anguilla vulgaris*) and *Leptocephalus morrisii* has been watched through its metamorphosis into the Conger Eel (*Conger vulgaris*).

If the contranantant spawning migration is against the S. W. monsoonal current, the ova and larvae will drift in a N. E. direction and those that enter the Straits of Malacca, for instance, would gradually approach the West coast of the Peninsula. Similarly, a spawning migration in the South China Sea during the N. E. monsoon would result in the larvae being carried along and dispersed along the East coast of the Peninsula.

As the larvae approach the coast they come within the influence of the tides and while continuing their progress with the monsoon current they are carried backwards and forwards by the daily ebb and flow of the tides.

¹ Meek, Migrations of Fish.

Their density causes them to sink lower in brackish water until they eventually find bottom in the shallow bays and estuaries and in this way are gradually dispersed all long the coast. Then a metamorphosis takes place and the feeble *Leptocephalus* is transformed into the active little fish which swims vigorously against the current and feeds incessantly and voraciously all the time.

In a recent report on the Fisheries of the Straits Settlements and Federated Malay States the writer drew attention to the Chinese fish-traps called *pompang* and other licensed fixed engines known as *ambai*, *langgai*, etc., of which there are several thousand between Penang and Port Swettenham. Though there are many kinds of these traps they all work on the same principle. In every case there is a wide V-shaped entrance terminating in a long funnel-shaped bag made of sacking or plaited split bamboos. The position of these traps is arranged with respect to the currents and tides so as to intercept the larvae and immature fish during their denatant drift to the shallows. Most of these traps float, and swing round with each tide so as to take toll both with the ebb and the flow.

An examination of the contents of these traps shews that in addition to immature fish, which any Malay fisherman will tell you are the fry of valuable food fish, the bulk of the catches are made up of feeble, attenuated, small-headed larval-like fishes which the Malays call **Bunga ayer** and to which they attach no value.

There can be little doubt that scientific investigation will prove that the **Bunga ayer** are valuable food-fish in the *Leptocephalus* stage.

This subject has been treated at some length because of its great economic importance and because the questions raised cannot be answered except by a specialist in marine biology.

Though myriads of larval and immature fish are caught daily for duck food, pig food and manure, and thousands of pikuls are exported as dried fish refuse, it has been argued, while admitting *ambai* catches are used mainly as pig food, that it appears a debatable point whether the flesh value thus produced is not as great as the extra fish value which might be caught if the fry killed by *ambai* were left undisturbed!

We cannot afford to allow such points to remain debatable.

Let us go on with the life history of the tiny fish which we left in the first stage of an active existence in the shallow waters near the coast. These shallows are the nurseries or recruiting grounds where the fry keep together in schools or shoals.

“After a period in relatively shallow water, the shoal migrates to deeper water. At first the migration is not to a great distance, but with growth the annual pulsation becomes greater and greater.

"The migration is not merely inshore and offshore, but is at the same time in a definite direction with respect to the coast.

"Thus the life of the fish is spent until in from three to six years at the most, the call of maturity comes. In response thereto a migration takes place which appears to be usually beyond the limits of the seasonal migrations of the school."¹

A few moments' consideration will enable one to realise that the life habits of every species of fish are subject to certain fixed laws. It is only a matter of systematic organised research to discover those laws and to apply the knowledge to the development of Malayan Fisheries.

We can learn what has been done in Canada, Great Britain and the United States, but this general learning must be supplemented by detailed local research. We must work out our own local tables.

There are, probably, no less than 2000 species of fish in Malayan waters. There are certainly not less than 500 species of economic importance, and if we take 250 species as being valuable Marine food fishes, some idea may be formed of the amount of research required before we shall be in a position to state definitely where a certain species may be found in full roe, where its spawning grounds are, where the recruiting grounds of its young are and when and where it travels during its seasonal migrations.

Information of this kind will enable our fish-hermen to catch fish in the best condition and in the greatest quantities and this is the information which the Fishery Departments of Canada and America give the fishermen, even to the extent of using aeroplanes, fitted with wireless, to locate shoals and disseminate information.

There is a great deal of knowledge, of which no use is being made, in the possession of many illiterate Malay fishermen, spread over wide areas, all along the coasts of Malaya. This knowledge should be collected and tabulated.

The Departments of Fisheries in Ceylon, Australia, India, the Netherlands Indies and the Philippines have published records dealing with the fishes which also inhabit our seas and, in consequence, the Fishery Officers and scientists have the benefit of a vast amount of scientific research work on which to build up local data.

Though the question of damage to our marine fisheries has evoked some attention during the past two years, it is doubtful whether serious thought has been given to the terrible damage done to the fresh water fisheries by mining silt. Engineers have fought for their roads and railways against the invading silt, but, to judge from official reports, no one has fought for the fisheries and the need for protection of the riverine rights of the people would appear to have passed unnoticed.

¹ Meek, Migrations of Fish.

Within the writer's memory the main rivers of the West coast were fine clear streams. The waters provided irrigation for the rice fields and contained quantities of fine edible fish. These rivers are now thick turbid streams carrying a heavy burden of slime and silt.

We have probably one hundred different species of Carp alone, besides dozens of species of Catfish and many fine fish belonging to the families OSPHROMENIDAE, NOTOPTERIDAE, etc., etc. Catfish can exist in slime and silt though it is questionable whether they can thrive, but Carp certainly require clear water to breed in.

One of our Carp the **Kélah** (*Barbus* sp.) has been described by Swettenham as the finest fresh water fish he ever ate in the East, and the **Kalui** (*Osphromenus olfar*) is so highly esteemed that several attempts have been made to introduce it into France, and it has been acclimatised in Mauritius, Australia and parts of India.

Tin mining is necessary and some pollution of the rivers is unavoidable, but there have been many cases where carelessly constructed dams have broken and a turbid flood of slime has been allowed to pour direct into the rivers for months while leisurely repairs are being made. Though much of the damage done in the past is irremediable, let us hope that a more general recognition of the value of the fresh water fisheries will result in a fair measure of protection in the future. There are still rivers which can be saved.

By saving our fresh water fisheries we shall save, incidentally, our rice-fields, for Rice and Fish in addition to being the two staple foods of the country are inseparable. When you destroy one you destroy the other.

Where you can grow rice you can catch fish and where you can no longer catch fish you cannot grow rice.

To explain: the mining silt which pours into the rivers gradually raises the bed of the stream and so causes a rise in the water table. A rise in the water table limits the area of drainable land, and drainage is as necessary to a rice field as irrigation. So the area which can be planted with rice becomes smaller and smaller until eventually the water table is so high that the river channel can no longer carry off storm water. The resultant floods deposit a layer of slime and silt on the rice fields and complete the work of destruction.

Fish cannot breed in the rivers polluted with slime and silt, so the Fisheries and rice fields perish simultaneously. In our policy of construction and development these facts should not be lost sight of.

There is yet another point which has received no attention and that concerns anadromous Marine fishes which enter rivers to spawn. Among these fishes the principal one is the Shad (**Téru-**

bok), which ascends the rivers to a considerable distance during the breeding season. It arrives on the coast in enormous shoals, and twenty eight years ago, as Skeat has recorded, they were invariably taken in full roe, when they are in the best condition.

Recent reports show that **Tērubok** have fallen off both in quantity and, as the writer knows from his own experience, in quality, those now taken being mostly spent fish in which state they are positively unwholesome.

These fish used to be taken in such numbers that the nets contained more than the boats could load. Within the past few years the writer has, on several occasions, picked up these fish by hand in a dying condition apparently choked by silt in their attempt to ascend the rivers. Failing to ascend the rivers the Shad must either spawn in the sea or in the polluted lower reaches and in either case the eggs perish.

Unfortunately, the migrations of the **Tērubok** do not, as far as the writer's experience goes, take it to the East coast of the Peninsula, so that, the **Tērubok** fishery of Malaya appears to be in danger of extinction.

This introduction would not be complete without some mention of the conditions under which the transport of fish from the source to the consumer takes place.

There is a general agreement that transport is bad. Many schemes have been evolved for ensuring rapid transport and reduced prices, but none of them have been put into practice and probably none are commercially practicable. A permanent scheme is required that can be built up by degrees; the writer has advocated in two reports the use of cold storage. While allowing that the expenditure will be great we should not lose sight of the fact that it will be a permanent and sound investment.

Let us consider the existing conditions first.

In a temperate climate fish will keep fresh for days. Here, near the Equator, fish caught in the morning are in an advanced state of decomposition before the evening. Decay is arrested by the use of ice. For instance, ice manufactured in Kuala Lumpur is taken by train to Port Swettenham and sold to small middlemen who go to sea and purchase from the fishermen. These middlemen are bound as a rule to sell the fish to the ice dealers, who again sell to other middlemen, who sell to the retailers in the markets. The result is that fish costing \$15 a pikul at sea cost \$80 a pikul in Kuala Lumpur, 30 miles away.

Ice melts rapidly in the trains, in the boats, and in the markets. A box of fish must therefore contain an enormous proportion of ice to allow for wastage, and the fish instead of being fresh, cold, and wholesome are in a swollen and sodden condition.

While these are the conditions under which fish are transported a few miles in this country, we are indebted to a single Cold Storage Company for the privilege of being able to purchase, if we can afford it, fish, meat, game, butter and fruit, imported in refrigerated chambers from Great Britain, the United States, Australia and China.

Briefly, it amounts to this. We can eat foreign fish and foreign fowl but not the fresh produce of Malaya. Hundreds of tons of prime fish are caught every year on the East coast, where the inexhaustible supplies of the China sea are available, but all this fish is dried for export for lack of cold storage transport, though much of it is caught within 24 hours steam of Singapore.

There can be little doubt that the whole future of the perishable food business in this country depends on cold storage, but there is no decided opinion as to the part that the State should take in the development of the trade.

It was realised many years ago, that for sanitary reasons the ordinary shop house was not a suitable place in which fresh meat, fish, etc., could be exposed for sale, and, in the Malay States, the sale of such perishable produce is confined entirely to the markets built by the State.

It would seem, therefore, to be but reasonable and logical for the State to go a step further, and instal cold storage in the markets, and to rent space to the retail dealers in the same way that stalls are rented.

The State owns the railways which run from the coast to the market towns and the installation of refrigerated vans on the railways would appear to be a natural development of a State enterprise, as it is in other countries with State Railways.

This disposes of the problem as far as the Colony and the West Coast States are concerned but the problem on the East coast is quite different.

The development of the States on the East coast has been retarded because they possess no natural ports and harbours which can be entered during the North East monsoon.

Though the deep sea can be fished all through the N. E. Monsoon and steamers run regularly up the East coast to Bangkok and Saigon, no fishing is done because the fishermen live on the mainland. A heavy sea breaks on the shallows and sandbanks which extend from the coast, and dangerous rollers break on the bars which guard the entrance to the rivers.

Further out, in twenty fathoms or so, the seas are regular, and conditions for fishing far better in every way than they are in a strong wind in the English Channel or in the North Sea.

We know that the sea off the coast of Pahang, Trengganu and Kelantan swarms with fish all the year round, and all that is necessary is a scheme for supplying the Western States and the Colony, where fish is now very scarce.

The writer advocates State enterprise in the establishment of cold storage depots on the islands, where there is always safe anchorage and shelter in smooth water.

There is a chain of these islands all the way up the East coast. An island with a cold storage depot will become a permanent fishing settlement. Rent would be paid by the fishermen for space in cold storage, to be collected when the fish is sold. So far State enterprise is advocated.

It would pay steamers, running from Bangkok, Saigon and China to Singapore, Port Swettenham and Penang, to call at these islands for fish, and those steamers not now fitted with refrigerating plant would instal it.

Schemes for ameliorating the lot of the fishermen by granting loans, etc., have not succeeded because no scheme protected the fishermen from the middlemen, but the depots which will be the Penny-banks of the fishermen, always ready to receive deposits, however small, until required, will render the fishermen independent of the middlemen.

For example, there would be nothing to prevent a group of Malay fishermen from consigning regular shipments of fish direct to a Malay retailer in the market.

Shipments would be so frequent that loans should be unnecessary, but allowing that loans were asked for, to start Malays working on a co-operative basis, as indicated above, there would be no risk in advancing money on the security of the stock of fish.

With State organised depots and State transport there would be a fair field for steam trawlers and steam drifters owned by Companies or individuals. The depots would receive the fish and save the trawlers a journey to port with every catch, and here again the middleman would be eliminated.

This work deals, very inadequately, with fishes only. Much could be written and will, no doubt, be written later about our Crabs, Prawns, Crayfish, Pearl oysters, Edible oysters, Scallops, Cockles, Corals and Sponges, but considerations of space prevent more than the briefest mention.

The writer has seen Pearl shell taken close to Singapore and has handled a pearl valued at £800 taken off the Kelantan coast.

Rock oysters grow well here, but as they take about three years to mature, and no native can resist taking them while still small, they are practically unknown in the markets.

Leases could be granted and oysters cultivated. Sponges too, can be cultivated. Commercial sponges can be grown from cut-

tings, like flowers, and are so grown in the Philippines, and there are yet other marine growths which can be cultivated in the gardens of the sea.

Few countries have the potential fishery advantages that we possess and have neglected hitherto.

Our position between the Indian Ocean and the China Sea is unique and not only gives us access to an unlimited area for deep sea fishing, but also accounts for the large number of species of fish.

From Kuala Perlis on the West to Kuala Tabar on the East we have a thousand miles of coast line; more than some nations possess.

As to the vital importance of fish in the diet of all dwellers in this tropical country there is no question: as to the existence of an enormous area of potential fishing grounds there can be no dispute, and as to the large variety and good edible quality of our fish there is ample proof. Can we doubt therefore, with the experience of other countries to guide us, and while remembering that the economic stability of every country depends on the price of the peoples' food, that our fisheries are capable of enormous expansion and can we doubt that an enlightened policy of exploitation and regulation combined with constant scientific investigation will render the Fisheries one of the great economic assets of Malaya?



Malayan Fishes.

PART I.

GIANT HERRINGS.

(*ELOPSIDAE*.)

This family contains but few species; the individuals however, abound in the tropical seas and are of great importance both as food and game fishes.

The **Bandang** or **Měnangin** (*Elops hawaiiensis*) is known to Americans in the Philippines & Hawaii as the Ten-pounder. It reaches a length of about 4 feet and is an edible fish of considerable value.

The **Bulan-bulan** (*Megalops cyprinoides*) is the Indo-Pacific Tarpon and is very closely related to the well known sporting fish of America. It is known in Australia as the Ox-Eye or Big Eyed Herring.

It has a very wide range extending from India to Australia and from East Africa to the Sandwich Islands. It is known to penetrate the rivers right up into fresh water and has been successfully cultivated in brackish or even fresh water.

From an edible standpoint it ranks very high. Its flesh is firm, well flavoured and possessed of good keeping qualities. It attains a length of 5 feet and when our fisheries are better known it may, as Stead has remarked, "turn out to be as great a sporting fish as the Tarpon."

THE FEATHER BACKS.

(*NOTOPTERIDAE*.)

The **Bělida** (*Notopterus notopterus*) is a fairly common fresh water fish which attains a length of well over three feet. The belly is said to be extremely rich and well flavoured but the back contains numerous small bones.

THE MILK FISH.

(*CHANIDAE*.)

The **Bandang** or **Jangas** (*Chanos chanos*) is the well known Milk-fish or White Mullet and is known as the Salmon-Herring in Australia. It is a sea and estuary fish and feeds on "sea moss," an alga (*Oedogonium*).

Its cultivation is a considerable industry in the Philippines, the Sandwich Islands and Java, and Day records that, in South Canara, Hyder Ali introduced it from the sea into tanks of fresh and brackish water where it still thrives.

Seale¹ gives a most interesting and valuable account of the cultivation of this fish, which is known by the Filipinos as the *Bangos*; I quote the following extracts:

“The Milk-fish is one of the most important commercial fishes in the Islands. It is raised chiefly in the fish ponds at Malabon and at other places near Manila and therefore can be secured at any time regardless of the weather.

“This fish is particularly adapted to pond culture being a vegetable feeder of rapid growth.

“The eggs are deposited in the sea. The young appear during the months of April, May, June and July. They are to be found in great numbers along the beaches and are captured by the natives and placed in large earthen jars full of water called *palyok*. They are then conveyed to the fish ponds, frequently a hundred miles distant.

“One of the jars contains about 2,500 young *Bangos*. About 60,000 are used to stock one pond of 1 hektare. As the fish grow they are thinned out by transfer to other ponds. Thirty-three per cent should reach marketable size and a yearling should measure half a metre.”

FOOD OF THE MILK FISH.

“If it is desired to cultivate the food alga, the water of the pond is allowed to drain off and the clay is exposed to the full power of the sun. The alga rapidly makes its appearance and a little water is then permitted to cover the bottom. This is gradually increased as the *Oedogonium* develops.

“The average value of the ponds about Manila Bay is probably 40 centavos per square metre, giving a total of more than 6,000,000 pesos for the pond value alone, which I am convinced is a conservative estimate.”

There should be no great difficulty in establishing a similar industry in Malaya and there are many mangrove areas on the West Coast of the Peninsula where series of ponds could be constructed.

There are many places where these fish are feeding on beds of sea moss and I saw millions of fry not far from Butterworth recently (March 21st) which a Javanese told me were **Anak jangas**. The Milk Fish attains a length of 5 feet. It is not often captured as it will not take a bait and will jump over a seine or drift net.

¹ “Fishery resources of the Philippines.”

HERRINGS.

(CLUPEIDAE.)

This is a very large and important family. The members range in size from the **Parang-parang** (*Chirocentrus dorab*) which is said to exceed a length of 12 feet to the **Bilis** (*Stolephorus tri*) which measures not more than 3 or 4 inches.

Although this family is of great commercial importance in Malaya, and the Herrings, Shad, Sprat, Sardines, White-bait and Anchovies belonging to it are highly esteemed for their flavour and food value by the Malays and all Eastern races, they are unknown to the great majority of European residents in this part of the world, with the exception of the **Bilis**, which is occasionally seen served as "White bait" or as a *sambal* with curries or in bottled form as Macassar Red fish.

From an economic point of view this family is second to none in importance and the fact that some of the most valuable kinds associate at certain periods in immense shoals accounts for the usefulness of the family as a food supply.

The following are the most important members of the herring family in our waters:

The **Parang-parang** (*Chirocentrus dorab*), the **Těrubok** (*Clupea* (*Alosa*) *macrura*), the **Sélangat** (*Dorosoma* spp.), the **Tamban** (*Clupea* (*Harengula*) spp.), (*Dussumieria* spp.) and (*Spratelloides* spp.), the **Bilis** (*Stolephorus* spp.) and the **Bulu ayam** (*Engraulis* spp.).

The **Parang-parang** is a very bony fish of excellent flavour and its capture by hand line provides a livelihood for several hundred Malays in Singapore alone.

Passengers by steamers proceeding through the Eastern entrance to Singapore roads will see a large number of small canoes in the deep water channel and will hear the noise of the rattles, which each Malay fisherman wields unceasingly. These rattles do not attract the fish, but keep the hand occupied and the fisherman on the "qui vive." The **Parang-parang** is not a greedy biter and does not stay in one place. He is a rapid swimming predacious fish who has no time for more than a snap as he darts through the water. Bites are usually few and far between and an inexperienced or somnolent fisherman would catch nothing. With an ever moving hand engaged with a rattle the fish is struck and hooked almost at the instant he bites.

The **Těrubok** is a Shad and is considerably larger than the ordinary herring. It is known to Europeans in India as the "Hilsa" or "Sable fish." Day says:

"They are excellent as food until they have deposited their ova, when they become thin and positively unwholesome. Their flavour has been compared to a combination of that of the salmon and herring: they are rather heavy of digestion."

The roe of the **Těrubok** (**Tělor těrubok**) is a highly valued delicacy and the fish are still caught in large quantities near Běng-kalis (Siak) and the roes dried and salted for export. Cantor writing in 1850, states that 40 years ago the Raja of Siak is said to have obtained an annual revenue of 72,000 guilders from the **Těrubok** roe industry alone.

The **Sělangat** of the southern part of the Peninsula is known as **Nandong** in Penang and Kedah, and as **Kěbasi** on the East coast. It is also known to the Filipinos as **Kabasi**. The English or American name for it is the Basling Shad and in Australia a member of the same sub-family is known as the Bony Bream or "Hair-back."

It is a bony little fish but very plentiful and cheap; it is dried and salted in enormous quantities both for local consumption and export.

The **Tamban** is one of our commonest fish and will eventually be one of our most important and valuable food fishes.

Dr. Cantor who wrote a catalogue of Malayan Fishes in 1850, says of the **Tamban bulat**: "It is highly valued for its delicate flavour and passes commonly as a 'sardine'": and of the **Tamban nipis** he writes, "They pass in the Settlements of the Straits under the denomination of 'Sardines' in imitation of which they are sometimes preserved in oil."

The Moros in the Philippines have the same name **Tamban** for this fish and Seale writes "Any of these sardines would compare favourably with the species put up in oil on the Pacific Coast."

We continue to import thousands of cases of sardines annually into the Straits and F. M. S., while our seas swarm with sardines.

It is popularly supposed that sardines are preserved in olive oil but I have more than a suspicion that refined coconut oil, sesame or gingelly oil, and other vegetable oils, which are largely exported from the East, return to us with the imported tinned herrings and sardines.

The **Bulu ayam** and **Billis** (Moro: **Dilis**) are anchovies and sprats.

Stead writing of the *Engraulis antipodum* of Australia, says, "For all practical and economic purposes there is no difference between our Anchovies and the famous fishes of that name in the Mediterranean Sea."

A glance at the systematic list of members of the herring family will show that we have at least 8 species of *Engraulis* and 3 species of *Stolephorus*.

QUEENSLAND-SMELT, ETC.

(SCOPELIDAE.)

One of the best known members of this family is the **Lumi** or **Luli** (*Harpodon nehereus*) which, when newly taken, is brilliantly phosphorescent all over the body: in a salted and dry condition it is the "Bombay-duck" so highly esteemed by Anglo-Indians.

It is quite common in our waters. I have seen many immature specimens in purse-nets, but full grown specimens are often taken in sunken basket-traps off Singapore in the neighbourhood of the Karimon Islands.

Day says "this fish is highly esteemed as food whether fresh or salted." It is curious that this fish which is a delicacy in India should be disregarded in Malaya. Malays do not care much for it, as the flesh is soft and flabby and they prefer firm and flaky meat.

The **Bělungkor** (*Saurida tumbil*) is known to Australians as the Queensland Smelt and is a fair edible fish, though somewhat dry.

The **Mudin** (*Saurus myops*) is a useful food fish.

"MILLIONS."

(CYPRINODONTIDAE.)

I have called these fishes "Millions" in the absence of any English name, because they belong to the same family as the fish known as "Millions" in Barbados. To this latter fish, which eats mosquito larvae, as our fish does, the immunity from fever, which the inhabitants of Barbados enjoy, is attributed. It is a tiny fish, very common in swamps and drains in Singapore and elsewhere.

CAT FISHES.

(SILURIDAE.)

This family which contains one fourth of the known fresh water fishes of the world is not represented in the British Isles, but in this region they are to be found wherever there is water and a muddy bottom, whether it be fresh, brackish or salt water.

Members of the family may be found in swamps, pools and roadside drains and many of the fresh water varieties will make considerable journeys overland to find new pools or streams. They are found in all our rivers and some species may be caught miles out in the sea. Some of the largest exceed 6 feet in length.

With the exception of the **Lele** and **Lembat** (CLARIIDAE spp.) which have no spines, all the members of this family are armed with spines.

The **Tapah** has a pair of pectoral spines only, but the **Sēm-bilang**, **Patin**, **Lawang**, **Pêdukang** and **Baung** have both pectoral spines and a dorsal spine.

A wound from these spines is extremely painful and the angler who captures one of these fish for the first time is advised to take a lesson from a Malay in the proper method of grasping them, which is very simple but worth knowing.

They are valuable food fishes and are in great demand among all Malays, Chinese and natives of India. Some species are considered to possess exceptionally nourishing qualities and are prescribed for patients recovering from illness.

They will live for hours out of water and can be transported for long distances.

The popularity of the Krian district of Perak among natives of India is due primarily to the rice fields and secondly to the fact that cat-fishes, as well as other fish, swarm in the rice fields and irrigation ditches.

A volume might be written on the Cat-fishes alone. One interesting characteristic is the great care they take of their eggs and young. The **Pêdukang**, for instance, lays very few but very large eggs which look like gelatine capsules and these they carry about in their mouths. As the **Pêdukang** are among the commonest estuarine fishes, any observer can obtain them during the breeding season and see for himself the egg in every stage of development, and in the final stage, immediately before hatching, the tiny fish is distinctly visible through the translucent envelope of the egg.

LOACHES AND CARP.

(*COBITIDAE* & *CYPRINIDAE*.)

Only two Loaches, the **Ikan pasir** and the **Lali**, are mentioned in this work, but judging from records of species in Java, Sumatra and Borneo, there should be at least 20 species.

The **Ikan pasir** (*Acanthopsis choirorhynchus*) is quite common in the Pahang river and is good eating.

The Carp family of fresh water fishes to which our Roach, Tench and Gudgeon belong is represented in our Malayan rivers by certainly not less than 100 species.

The very incomplete list in this book gives some 28 Malay synonyms only. There is an interesting hobby and good sport with a fly and spinner awaiting any planter or prospector who lives near the upper reaches of any of the rivers in the Peninsula. An oil drum can easily be converted into a specimen tank in which rare fish may be preserved in spirit. The Directors of the F. M. S. and S. S. Museums would be only too glad, I feel sure, to mount and display specimens, and as the field is practically untouched the collector has more than a sporting chance of discovering and perhaps giving his name to a new species.

During the breeding season, the males of many species assume a more brilliant livery, or develop excrescences and tubercles on various parts of the head, especially on the snout, or also on the body and fins.

The common Carp of Europe is said to have been introduced from China early in the seventeenth century. The Chinese continue to import Carp into Malaya and to grow them in stock-ponds. The imported Carp are the **Tiam** (Chinese) (*Labeo moli-torella*), the **Ling** (Chinese) (*Cyprinus carpio*), and the **Hwan** (Chinese) (*Ctenopharyngodon idellus*).

The ova are shipped from China in large jars full of fresh water and the contents of the jars are regularly and vigorously stirred with a stick or paddle during the voyage, to oxygenize the eggs, and by the time the jars arrive in this country they contain thousands of fry.

These fish are very popular among the Chinese and fetch high prices in the markets. They attain a length of three feet or more and a weight of perhaps 20 to 25 pounds. The utilitarian owners feed them on food of such a disgusting nature, principally excreta, that I can say nothing as to their edible qualities, as I have never felt any wish to taste them.

I imported some many years ago and intended to stock a pond in Kuala Pilah but unfortunately the ship was placed in quarantine and as their period of confinement in jars is limited, all the fry perished. These imported Carp have not, so far as I know, been bred in this country and it is more than likely that they require fresh running streams for the natural development of their ova.

The question of stocking some of our streams may be worth consideration, but I doubt it, as we have so many indigenous Carp. In this connection it is well to remember that these Carp which have been artificially bred for centuries, have yielded numerous examples of hybridism. I have read in an American magazine of a sportsman who for lack of other bait used aquarium gold fish (Carp) very successfully as live bait. He kept a stock of them in a fountain where they interbred with small species of American Carp with the most extraordinary results.

Professors Max Weber and de Beaufort write of the *Cyprinus*, "Distribution: Fresh water of temperate parts of Asia and Europe, from where introduced in many parts of the world and changed into many varieties."

We have many species of *Barbus* including the famous sporting fish the Mahseer of India, our **Tēmoleh** (*Barbus mosal*). The **Kérai** (*Barbus neilli*) is said by Day to attain a weight of 50 or 60 pounds.

The Catla (*Catla burchanani*) of India, Burma and Siam has not yet been identified in our rivers and as it attains a length of at least 6 feet and a corresponding weight, I suggest the quest of the Catla, as a sound sporting proposition, to the anglers of Kuala Lipis, and other up country stations, with a good chance of success.

EELS.

ANGUILLIDAE, MURAENIDAE, CONGRIDAE, ETC.

The Eels are placed on the Order APONES which has several families and not less than 30 species, represented in Malayan waters. Eels are more popular perhaps with the Chinese than with any other race in Malaya. The Congers (**Malong**) are quite common and find a ready sale in the markets.

One eel (*Thyrsoidea macrura*) known as the **Pampan** or **Pompa** by the Javanese is probably the largest eel in the world and exceeds 10 feet in length. It has been found in the shallow seas and estuaries, and even rivers, of Sumatra, New Guinea, Natal, India, Ceylon and Queensland, but has not yet been recorded by local scientists.

SWAMP-EELS.

SYMBRANCHIDAE.

The **Bělut** (*Monopterus albus*) is confined to the fresh or brackish waters of Burma, the Malay Archipelago and China.

“This eel is numerous at Chusan, in streamlets, canals, and estuaries. As it is a favourite article of food it is kept by the inhabitants of Chusan in large jars, with fresh water. But it is capable of living a considerable time out of water. It is of voracious habits, feeding on smaller fishes, and it takes hooks baited with earthworms.” (Cantor).

SEA-HORSES AND SKELETON-FISHES.

(SYNGNATHIDAE & AMPHISILIDAE.)

The little Sea-Horse, **Kuda-kuda laut** (*Hippocampus* spp.), which is so like the conventional knight of the chess-board, is a familiar object in most Museums. It has, of course, no economic value.

A peculiarity of this curious family is that “the males are provided with a pouch (in some species there is only a broad groove) in which the eggs are hatched after being deposited by the female.

“The males take full charge of them, and the young remain in the pouch usually for a short time, after being hatched.” (Stead).

The Skeleton-Fish, as I have called it, *Amphisile scutata*, is known to Malays as **Ikan kĕring**. It is only a few inches long and, as it has practically no flesh on its bony carcase, it has no edible value.

GAR-FISH, GAR-PIKE AND FLYING-FISH.

(SCOMBRESOCIDAE.)

Most of the members of this family, of which about 200 species are known, are marine: some are carnivorous, and others mainly herbivorous, feeding on green algae. Nearly all are in the habit of making great leaps out of the water, this tendency culminating in the Flying-fish (*Exocoetus*), and there is every passage between the small pectoral fin of a Gar-fish and the swallow-like "wings" of the most developed Flying-fish.

They are excellent edible fish and the **Todak** (*Belone* spp.) which have green bones are nevertheless very palatable and useful food fish. There is a more or less prevalent idea that fishes with green bones are poisonous; this is one of the popular fallacies which no facts or arguments will ever overcome.

The **Puput** or **Jénjulong** (*Hemirhamphus* spp.) or Half-beak can be easily identified by the fact that the lower jaw only is prolonged. Both jaws of the **Todak** are prolonged into a long slender beak.

The **Todak** though common is a very shy and wily fish. It gives a boat a wide berth and though a rapid swimming and predacious fish it disdains ordinary lures and baits. The Malay fishermen, however, circumvent them by fishing with a kite from which dangles a length of fine line terminating in a noose. The bait is attached to the noose. Given a breeze sufficient to keep the kite flying, quite good sport is obtained, perhaps a dozen fish or more in an hour. The **Todak** makes a fierce dash at the bait, the noose tightens round the upper jaw, and the recurved teeth prevent the noose from slipping.

The **Puput** is also a shy biter when it sees its would-be captor at the other end of a rod or line, and the Malays consequently use short baited lines attached to floats (*Pëlontang puput*) which they send adrift and pick up when the movements of the float shew that a fish is firmly hooked.

Flying-fishes are very good eating but not often obtained. I have picked them up on the deck after a squally night at sea.

SAND-SMELTS.

(ATHERINIDAE.)

These pretty little fishes, called **Rennyau** or **Paku** in Malay, are common all along our coasts and also frequent estuaries and tidal river. Seale¹ writes:

"It is almost impossible to land at any wharf or to go ashore on any beach without seeing these little fish in countless numbers. They usually grow to a length of from 10 to 12 centimeters. They have a greenish tint on the back and a bright silvery band on the sides.

¹ Fishery resources of the Philippines.

"There are five or six different species.

"The most common species is possibly *Atherina temminckii* (Bleeker).

"They are known as 'pescados del rey' or fishes of the king, among the Spaniards.

"They are greatly valued as food. The young are termed white bait. The method of catching is usually by seine or corral. A profitable industry could be built up by preparing these fish in a good sauce, by pickling them with spices, or by drying. They abound at all seasons."

GREY MULLET.

(*MUGILIDAE*.)

The Grey Mullet are a widely distributed and very important family. They inhabit shallow water in the seas, estuaries and rivers and none are known to occur in very deep water.

Their habit of keeping to the shallows, in large shoals, renders their capture, in enclosures, which dry out at low tide, and in mullet nets, an easy matter. As there are no restrictions as to size and no close season, mullet are getting scarcer every year in Malayan waters.

Grey Mullet feed, more or less, on the organic matter found in mud and they are peculiar among fish in that they have a true gizzard, lined with a thick horny epithelium.

Mullet are very common and highly appreciated in Australia. Stead¹ writes:

"During a recent year in New South Wales alone 45,000 baskets of Mullet—principally Sea Mullet—were received for disposal at the various fish markets. The average basket of Mullet contains about 75 pounds weight; and, if we calculate the fish at an average of one pound weight each, we find that we have the imposing total of 3,375,000 individuals."

In New South Wales Mullet may only be netted at certain times and at certain places, and there is a legal limit as to size as with all valuable edible fish in that Colony.

The breeding season with most of our Mullet appears to be between November and February during the N. E. monsoon.

At this time I have seen the **Anding** in myriads in the surf, near the mouth of the Trengganu river and other rivers on the East coast. A quantity of yellowish foam and scum is brought down by the rivers which are usually in full spate in November and December, and this foam either contains food or provides a suitable shade and shelter for the ova of the Mullet which are, I believe, pelagic or floating eggs. A day of steady incessant tropical rain during the N. E. monsoon is the day above all others to which all

¹ Fishes of Australia.

Trengganu Malays, male and female, look forward. When the rivers are in full flood, the sun obscured, the N. E. monsoon blowing half a gale, the surf thundering on the beach and full of yellow yeasty foam, then you will see all the Malay ladies trooping out in their best silk coats and sarongs, and all the old blades and young bloods are in attendance.

They are all out for the day to enjoy themselves and to catch mullet and the more it rains and blows the better they like it, the ladies, perhaps, because their vivid silken raiment looks best when it is wet, or may be it fits their figures better so, and the men, perhaps, because they will catch more mullet!

These ladies have designed and made their own costumes. Raw Chinese silk has been teased, wound and spun; fast dyes of vivid colours, orange, pink, vermilion, green; every colour and every shade have been prepared from roots, bark and leaves, and the garments have been woven in intricate designs, tartans, checks, watered silks and shot silks; a creative art which has been lost on the West, and will soon be lost on the East Coast, in these days of cheap imitation silks and aniline dyes. But let us get back to the mullet and the rain.

And the more it rains the fresher keep the flowers in the ladies' hair. These ladies wear no hats and there are no collars, dragged skirts or squelchy boots in this picnic party.

Let us again to the mullet. Now this catching of mullet is an affair of casting-nets and he who catches the most mullet is some ace. It is not a simple poaching trick of slinging a net over a sleepy fish in a pool, but quite a different business, I assure you.

The nets are made of the finest and strongest cotton, water-proofed in white of egg which renders them to the touch, for a season, as though they were made of the finest gut or sinews. The small net or *jala anding* when thrown covers perhaps 100 square feet of surface and it is weighted with little chains of pure tin. The light cord attached to the thrower's wrist is usually 30 feet long and the net is often thrown so as to drop fully expanded at the full extent of the cord, and that throw is in the teeth of a North-east gale.

Each fisherman has perhaps two or three such nets and, in reserve, a much larger and stronger casting net for the **Pělong** which is the giant of all our mullet.

Keeping far back on the sandy beach, the men follow the shore line until mullet (**Anding**) are seen, and, to the novice, it is a difficult matter to see them. But there they are, and when you know what to look for, in the smother and foam, you will notice little black heads, in hundreds, between the breakers. Now these **Anding** are the shyest fish that swim. A wave of the hand and they have disappeared to pop up again at a distance further seaward, where no man can hope to reach them.

This, then, is the manner of their capture. There will come a moment when a great wave, like a wall, hurls itself on the beach. In fact these waves do it all the time! However, there is measure of two or three moments and no more when that wave stands like a wall between you and the fish, and the fish forget your existence. In that brief time your caster of the mullet net sprints down to the very verge of the breaking wave and up to or over his knees in the water; the net truly held and truly swung, with a long pendulum swing, clears the crest of the approaching wave and falls fairly on the group of mullet concealed in the hollow beyond, and in this way perhaps he may be fortunate enough to take one or two hundred fish in one cast. But you will serve a long apprenticeship, and will, when learning, throw half a hundred times and have no mullet

When the sport is in full swing perhaps 10 or 20 men dart simultaneously down the beach and as many nets shoot out and over the waves. Suddenly there comes a wild yell of excitement. **Pělong** which have been disturbed or enclosed in the **Anding** nets leap several feet in the air and break their way through the nets. Silvery six-pounders and even larger fish instinctively jump when their brothers jump. There is a rush up the beach and a race back to the breakers with the **Pělong** nets. The nets are thrown at random (*tebar rambang*) in every direction. There may be a shoal of **Pělong** and, if so, some excitement I promise you.

Once a **Pělong** sees the net over him, he makes one upward dash to the apex of the net. The fisherman hurls himself at the fish and must grasp him then or not at all, for the next powerful dash for liberty takes the fish down to the bottom and he is under the chains and out of the net before you can wink. Out of your depth in a strong surf with a couple of lusty **Pělong** in your arms and a smother of net, chain and cord about you, you come to the conclusion that life was never more worth living and that if you are off to kingdom come you will take the **Pělong** with you.

As I write, at Tanjong Katong, Singapore, I can see some Boyanese, syces probably, with baby casting nets catching shrimps, sprats and baby fish in a sea like glass; a miserable messy business. The real gladiators of the casting net are to be found only on the East coast.

Our Mullet (**Bělanak**) include the **Jěmpul** (*Mugil planiceps*) which attains at least a foot and a half in length.

The **Tamok** (*M. waigiensis*), according to Day, attains at least 3 feet in length. The **Anding** and **Kědéra**, which grow to about a foot and a half, are excellent eating.

The **Bělanak tamok** (*M. waigiensis*) is known in Australia as the Diamond Scaled Mullet. It attains a weight of several pounds and is of a pretty silvery colour, each scale being prettily margined with black.

THREAD FINS.

(POLYNEMIDAE.)

These fish are all excellent as food and from some rough isinglass or fish sounds are obtained.

One of our **Kurau** (*Polynemus paradiseus*) is the Tupsi fish of India and the Mango fish of Burma. This small fish which attains a length of about 9 inches only, is considered a great luxury both in India and Burma. It has very long filaments, longer than itself, proceeding from each side, just below the pectoral fins.

The common **Kurau** (*P. indicus*) attains a length of 4 feet and about 20 lbs. in weight.

The **Kurau janggut** (*P. tetradactylus*), which is also known on the Queensland coast by the very unsuitable name "Cooktown Salmon," grows to a very large size.

Day quotes Hamilton Buchanan as observing "I have been assured by a creditable native that he saw one which was a load for six men and which certainly therefore exceeded in weight 320 pounds avoirdupois."

BARRACUDAS.

(SPHYRAENIDAE.)

The **Alu-Alu** are carnivorous and highly voracious fishes which give good sport with a trolling bait or spinner.

Cantor mentions two species only, *S. jello* and *S. obtusata*, as inhabiting the seas off Penang and Singapore but I have a snapshot of one taken by me off Pahang which does not agree with Day's plates of either of these fish and is I think *S. novae-hollandiae*. It is more than probable that other species will be discovered in these waters.

A well known American game fisherman from Honolulu who visited Singapore recently recognised the fish both in the Singapore market and from the photograph as the **Alu-alu** of Hawaii.

Roughley¹ writes of *Sphyraena novae-hollandiae*,

"The Short-finned Pike is an edible fish of very considerable value and is deservedly popular, though it has probably never been tasted by a very great portion of the community.

"It forms one of a too numerous collection of very valuable table fishes which are scarcely utilised, owing to the unenterprising methods adopted in their capture."

¹ Fishes of Australia.

POMFRETS.

(STROMATEIDAE.)

The **Bawal** is one of our well known and most popular edible fish and takes a high place among our very best food fishes.

Of the three species known, the **Bawal chermin** (*Stromateus atous*) is most esteemed, though all are good.

The **Bawal puteh** and **Bawal itam** (*S. cinereus*) are identical, the fish getting darker as it reaches maturity. A shoal of immature fishes of this species seen on a calm, clear night is a most beautiful sight. In the reflected light of the moon, they look like tiny silvery stars. The Malays call them **Bawal bintang**. When mature they are a darker gray colour.

The mature **Bawal itam** are very much better eating than the immature **Bawal puteh**.

The **Bawal tambak** (*S. niger*) is the largest of the family. It grows to a length of two feet and is excellent eating.

These fishes are considered by fishermen the stupidest fish that swim. They have a curious sheep-like habit of huddling together and are also afflicted with a kind of ovine curiosity. They will follow anything that excites their curiosity such as a boat and this habit leads to their capture.

When a shoal is observed the fishermen manoeuvre their boat so as to attract the interest of the fish until they are all following in a compact flock. At the same time the boat is taking a circular course and the net is being shot very quietly. At the right moment all the fishermen raise a tremendous din with clappers and at the same time splash the water with their paddles. The shoal does a perfect right-about turn and each fish dashes headlong into the net.

MURREL.

(OPHIOCEPHALIDAE.)

These fresh-water fishes are provided with a respiratory organ on each side, above and communicating with the gills, which enables them to breath atmospheric air.

They can live for long periods out of water and travel over the land from one piece of water to another. They are useful food fish and well adapted for pond culture provided that the pond is well stocked with the small fish and frogs on which they feed, but if the food runs short, they will go elsewhere.

Day writes, "Jugglers both in India and China exhibit these fishes walking on the land, and children amuse themselves by making them crawl along."

The young as a rule are of a more or less orange or scarlet colour.

These fishes appear to be monogamous, some breeding in grassy swamps or the edges of tanks and others in holes in the river banks.

They construct nests amongst the water-weeds where the ova are deposited. When very young the fry of all species, **Aruan**, **Toman**, **Bujok**, etc., keep with, and are defended by, their parents, but as soon as they are sufficiently strong to capture prey for themselves, they are driven away to seek their own subsistence: those which are too obstinate to leave being eaten by their progenitors.

The Malays have a saying *Bagai toman makan anak*, "Like the **Toman** fish which eats its own young," which is applied to persons in high places who misuse their powers, oppressing those whom they should protect.

The **Aruan** and **Toman** will readily take a bait, especially a frog, and are said to rise to the salmon fly. The largest run well over 3 feet in length.

They are caught in great quantities in the Krian irrigation reservoir at Bukit Merah and sent alive in tubs all over the F. M. S.

NANNYGAI.

(*BERYCIDAE*.)

The **Sěbėkah karang** (*Myripristis murdjan*) is a small fish of no particular economic importance.

The *BERYCIDAE*, of which there are about 70 species, live, mostly at great depths, in the seas all over the world.

The "Nannygai" of Australia, which belongs to this family, is highly esteemed on account of its delicate flavour and firm white flesh. Roughley writes,

"Until recently the supply of 'Nannygai' to the market has been an intermittent one, occasional specimens only being found there.

"The trawlers have now quite altered this and large quantities are being received from them daily, with the result that it is one of the commonest fish seen in the market.

"Hundreds of people visiting there in search of trawled fish are now seeing the 'Nannygai' for the first time."

I suggest that the capture of the "Nannygai" and other, hitherto unrecorded, species of good edible deep water fish, by means of a commercial steam trawler, is well within the region of possibility. We have, as a perusal of this book will shew, many fish in our waters which range as far as Australia but no engines or methods of capture are utilized in our waters which take bottom feeding fishes in depths of 50 fathoms.

Roughley writes, "The most prolific grounds are found to be between 50 and 60 fathoms in depth," and again, "Many species, which before the advent of trawling were very rarely seen in the markets and which were in fact considered by many to occur only in small numbers on our coast, could now be counted in thousands."

KNIGHT-FISHES.

(*MONOCENTRIDAE*.)

The **Sětonggang** (*Monocentris japonicus*) is a curious little fish with enormous bony scales and two long ventral spines. It has no edible value.

"BULL'S-EYES."

(*PEMPHERIDAE*.)

I have no personal knowledge of these fishes and, as they are never captured in numbers, they are unimportant from an economic standpoint.

"DRUMMERS."

(*KYPHOSIDAE*.)

The **Tėlan rumput** (*Kyphosus cinerascens*) as its name explains is a herbivorous fish, subsisting chiefly on "sea-grass" and "sea-moss." It is a congener of the Drummer of New South Wales (*Kyphosus sydneyanus*) and belongs to the same family as that excellent food and sporting fish known in New South Wales as the Blackfish.

DUSKY-PERCH.

(*LOBOTIDAE*.)

The **Pėchah pėriok** (*Lobotes surinamensis*) is a large perch-like estuary fish which reaches a length of 3 or 4 feet and a weight of 25 to 30 pounds. It is known in Australia as the Dusky Perch.

According to Boulenger's classification, the family contains two genera, *Lobotes* and *Datnioides*, each with two species, and though the two species of *Datnioides* are known to occur in the estuaries and rivers of the Malay Peninsula, they have not yet, as far as I am aware, been identified under a Malay name.

The **Pėchah pėriok** is an excellent food fish and it readily takes a fish bait.

It has a very wide range, being found in Queensland, the West Indies, on the east coast of the United States of America, as far North as Cape Cod, in the Mediterranean Sea, India and China.

BLOW-PIPE FISHES.

(TOXOTIDÆ.)

The **Ikan sumpit** or **Sumpit-sumpit** (*Sumpitan*, a blow-pipe) is so named from its method, unique among fishes, of shooting water from its mouth at insects which it perceives close to the surface.

They are very common estuary fishes congregating under piers, fallen trees and branches, where they may be both seen and heard spitting at flies and similar small game, which they knock down with surprising accuracy.

In the second edition of Day's Fishes, this shooting habit is erroneously attributed to another fish, *Chelmo rostratus*, which is also known to the Malays as **Ikan sumpit**, and a note under *Toxotes* reads, "It is stated in some works that these wide mouthed fishes shoot insects with a drop of water. The action is one which the mouths of these fishes appear incapable of effecting."

There is ample evidence, however, to prove that *Toxotes* do shoot, and though their mouths are large it will be noticed that they have the projecting lower jaw of the true cuspidore artist.

Chelmo rostratus, on the other hand, though it has pipe-like projecting jaws, is a fish which is found at sea in the neighbourhood of coral reefs where flies and insects must be rare.

It owes its name "**Sumpit-sumpit**" to the fact that, after capture, it spurts water through its mouth.

Toxotes chatareus grows to a length of about one foot and I have taken several with a rod when fishing with prawn bait for **Siakap**, between half and three quarters of a pound. They are quite good eating.

FRESH-WATER PERCHES.

(NANDIDÆ.)

I have no information regarding the **Kěpau** or **Patong** (*Catopra fasciata*) which is the sole member of this family recorded in our waters under a Malay name.

SEA-PERCHES.

(SERRANIDÆ.)

The very incomplete list in this book gives some 45 species of Sea-perches. The family is a very large and most important one, containing as it does, not only some of our largest fish but also many of our best edible fish including the **Ikan merah**. Nearly all the members of this family are carnivorous fishes which take a bait readily.

I propose to mention only a few of the most important ones.

The **Siakap** or **Kakap** (*Lates calcarifer*) is also found in the seas and estuaries of India where it is known to Europeans as the "Cock-up" and its range extends to Queensland and Western Australia, where it is generally known as the "Barramundi."

It is a fine sporting fish and runs to a considerable size. The largest that I have heard of was taken in the Bay of Bengal by the Government trawler "Golden Crown" and weighed 580 pounds. Before I read of this fish I used to be satisfied with 40 or 50 pounders.

The **Kĕrapu** (*Epinephelus* spp.) are very well represented in our waters and are fine edible fish. Some species lose their bright colouring soon after they are caught and have a dull mottled appearance when exhibited in the fish market which would not attract a purchaser unacquainted with the fish. From an edible point of view the **Kĕrapu** differs little from the **Ikan merah**, the latter fish owing most of its popularity to its colour.

Kĕrapu from 50 to 70 pounds in weight are occasionally seen in the markets and the **Kĕrĕtang** (*E. pantherinus*) is commonly seen up to two or three hundred pounds in weight. Any Malay fisherman will tell you of a **Kĕrĕtang** of fabulous size which he hooked and fought for hours, being worsted in the end because his boat and gear were too light to make any impression on the fish.

Very large ones are occasionally taken in fishing stakes (*Kelong*) and I have heard that the captors, on these occasions, tickle the monster until they get it quiet and then pass a strong rotan through its gills by which it is finally secured and hauled up.

The largest **Kĕrĕtang** I ever saw was taken by a Malay and myself. We had to sink our boat after we had fought the fish to a finish in order to load it; the weight, for there were no means of weighing it, was estimated at 6 piculs, i.e. round about 800 pounds.

This fish appears to me to be identical with the Queensland Groper illustrated in Stead's "Edible Fishes of New South Wales" under the name *Promicrops itaiara*.

I have read that this fish derives its name Groper from its habit of groping about the rocks but I suggest that the name originated in Malaya or India. The Tamil name is **Kurrapu**. The Malay name is **Kĕrapu**. The Brunei Malays know it as **Kurapa**. In the Philippines it is known to the Filipinos as **Garropa** from which the transition to Groper or Grouper is a slight one.

One of our **Kĕrapu** (*E. tauvina*) is known in Australia under the names Brown-spotted Hind and Black-spotted Rock-cod, and of this fish Roughley writes, "It is of fine edible quality and grows to a length of at least four feet."

In America members of this family are known as Sea-Bass.

The "Snappers" include our **Ikan merah** which is known as **Jěnehak** in Penang (*Lutianus* spp.). There are perhaps more than 20 species in local waters of which the list in this book gives 13 only.

Two or three Snappers are of a brownish colour, but the colouring of the rest of this brilliant family ranges between crimson, scarlet and golden, while some have violet, purple and blue bands.

They are all good edible fish and no swagger dinner in Singapore is supposed to be complete unless **Ikan merah** appears on the menu. Their popularity makes them expensive.

"WHITINGS."

(*SILLAGINIDÆ*.)

The **Bulus-bulus** (*Sillago* spp.) is one of our common market fish and can be obtained all the year round, though never in large quantities.

The *Sillago sihama* is known in Madras as "Whiting" and *Sillago maculata* is called the Trumpeter Whiting in Queensland and New South Wales where it is greatly valued for its excellence as a food fish.

They frequent shallow water and sandy bottoms where they feed on small crustaceans, worms, sand hoppers, etc. There is probably no cleaner feeding fish than the Whiting, a fact which perhaps accounts in some measure for its delicate flavour and wholesomeness.

Both our varieties, whether adult or young, are very shy and instantly bury themselves in the sand on the appearance of any danger. Even a passing dark cloud leads to their immediate disappearance into the sand whence they emerge a few moments later.

Roughley writing of another species, which has the same habit says:

"In the capture of this fish the hauling net is principally used. It displays considerable resource in evading the net, giving at times much trouble to the fisherman. As it is hauled near the shore, many fish, perceiving that they have been trapped, quickly burrow into the sand. Were not the fisherman alert to this cunning method of evasion, a large number of fish would be lost, but when it is known that the haul consists, in the main, of Whiting, they carefully tramp over the sand enclosed by the net and upon feeling any movement beneath their feet, quickly grab the concealed fish."

Whiting fishing is perhaps the nearest thing to trout fishing that the sea-angler can obtain. A light rod, fine tackle and small hooks are required and the sand flats should be fished on the flood tide. The bait should be cast as far as possible. The fish will be taken in water only ankle deep and the best bait are prawns, small bivalves, **Rěmis**, **Kěpah**, etc., which are found on sandy beaches, and beach worms, **Pumpun sarong** and **Pumpun darat**.

JEW FISHES.

(SCIAENIDAE.)

About 150 species of this family are found in various parts of the world. Nearly all are of economic value, some being highly so, and many of them reach a very large size.

The **Těmběreh** (*Sciaena diacanthus*) is one of the commonest coast and estuary fishes and perhaps the largest member of this family in our waters. It attains a length of at least 5 feet.

The **Gělama** (*Otolithus* spp.) are among our commonest fishes. They travel in shoals numbering many thousands and are taken in deep water hauling nets (*Pukat pětaram*) by Trengganu and Kelantan fishermen. These fish are dried and salted on the East coast and thousands of pikuls are exported annually.

The **Gělama** will take a bait but are hardly worth fishing for as they give no sport and are insipid table fish even when quite fresh. As "**ikan kěring**" with curry they are quite good.

"SILVER-BREAM."

(GERRIDAE.)

These are small fish inhabiting all tropical seas and entering estuaries.

According to Day these fishes are eaten by the indigent classes in India being little esteemed when fresh, but as they salt and dry well, large numbers are prepared in this manner for use.

The family contains about sixty species of which only six are mentioned in this work. Some 15 species are found in Australia and 23 in Indian waters. They rarely exceed a length of ten inches: nearly all have a plain silvery coloration.

In America, the fishes of this family are known as "**Mojarras**."

The **Kapas-kapas** (*Gerres* sp.) will take a bait, preferably prawns or beach worms, and may be caught in the vicinity of fishing stakes (*Kelong*) as in and also near reefs. When freshly caught it will be found quite a pleasant table fish with comparatively few bones.

SELEMAH.

(LACTARIIDAE.)

The **Sělěmah** is the sole member of this family and is not a fish of much economic importance. It grows to a length of about 10 inches, and is eaten by the natives either fresh or salted but is said to be insipid.

It appears in Malabar in shoals during the months of February and March.

They enter the Straits during the N. E. monsoon but not in large numbers and I am informed by Malay fishermen that they rarely take more than half a dozen on any one day.

They do not take a bait but a few find their way into nets and traps.

GRUNTERS.

(PRISTIPOMATIDAE.)

This family contains about 130 species belonging to four genera, of which three genera inhabit our waters, viz. *Pristipoma* (**Gērut-gērut**), *Diagramma* (**Tēbal bibir**) and *Pentapus* (**Sēlinching**).

The **Gērut-gērut** are good food fishes and take a bait readily. They are fond of back waters and one species (*P. guoraca*) is said to have been captured in fresh water.

I have taken several in brackish water and found them, when freshly cooked, excellent eating.

Our largest species (*P. hasta*) attains a length of about 18 inches and is known in Australian waters as the Queensland Trumpeter. Of this fish Stead writes, "The Australian home of this magnificent food-fish is principally along the coast of Queensland where it is well and favourably known."

The names **Gērut-gērut** and Trumpeter are descriptive of the grunting noise the fish makes after capture.

The **Tēbal bibir** are also good edible fish and attain a length of two feet or more.

I have no personal knowledge of the **Sēlinching** (*Pentapus*) and place it here from a description supplied to me, together with a pocket-kodak snap-shot which does not display the fish very well.

SEA-BREAMS.

(SPARIDAE.)

This, again, is an important family which includes many varieties of valuable food fishes. Some are carnivorous.

Following Dr. G. A. Boulenger's classification, the principal genera found in Malayan waters are *Scolopsis*, *Synagris*, *Caesio*, *Crenidens*, *Lethrinus* and *Sparus*.

Of the **Gērētak lantei** (*Scolopsis* spp.) of Singapore, so called from the parallel bands which distinguish most species, I can say little. The only species of *Scolopsis* with which I am familiar are the **Anjang-anjang** and **Kērisi bali**, which are occasionally taken when fishing for **Kērisi**.

The **Kērisi** (*Synagris* spp.) are beautiful little fish of a roseate hue with yellow and silvery bands. They are very common all up the east coast where they can be taken with a line, practically anywhere, in fairly deep water on a sandy bottom.

They average perhaps five or six to the pound but I have taken them up to a pound or more in 30 fathoms near Tioman Island. **Kērisi** fishing is, or used to be, the favourite out-door sport of the Malay Princesses of Pahang, and during the S. W. Monsoon regular expeditions were made to the **Kērisi** grounds and

the little fish would be hauled in until the boats were deep in the water and the Royal ladies exhausted. At the right season, there are few more delicate flavoured fish than the **Kĕrisi** and they remind one of really good Whiting.

But they must be absolutely fresh and caught on the right ground; if out of season or stale, **Kĕrisi** have an unpleasant tang about them.

The **Dĕlah** (*Caesio* spp.) are small but good eating, the best being, perhaps, *C. pinjalu* which is also known as **Ikan merah china** and is in no way inferior to the **Ikan merah** as a table delicacy.

Of the genus *Sparus*, three species are mentioned in this book, one of which, the **Bĕras-bĕras** (*S. sarba*), is the Tarwhine of Queensland and New South Wales, where it is considered a good edible fish.

It is not to be compared however, either from a sporting or an edible point of view, with its congener the Black Bream (*S. australis*), which has not been recorded as inhabiting Malayan waters.

The **Asoh-asoh** (*Lethrinus nebulosus*) is another useful fish in this family. The inside of its mouth is orange coloured as is that of its relative the Yellow-mouthed Snapper (*L. chrysostomus*) of Australia.

RED MULLET.

(MULLIDAE.)

Members of this family are known as Red Mullets in Great Britain and as "Goat-fishes" or "Surmullets" in America.

The British species are *Mullus barbatus* and *M. surmuletus*, remarkable for their beautiful pink or red colour, and much valued on the market, although no longer held in the high estimation for which they were noted by the Romans.

Biji nangka or **Lĕbai** are the Malayan generic names of our local members of this family and are descriptive. The **Biji nangka** (Jack-fruit seed) is yellow and has a filamentous process similar to the barbel of the Red Mullet; a **Lĕbai** is a Malay of exceptional pious habit, and it will be noticed that he almost invariably sports a beard consisting, as a rule, of about two or three long hairs, and his fellow countrymen have hit off the resemblance to the fish, which has two long barbels dependent from the lower jaw.

The **Ikan lĕbai** are remarkably beautiful fishes and their brilliant colouring contrasts somewhat with the solemn aspect of the head, which is, perhaps, an additional reason for the Malay nickname.

One of our local species (*Upeneus tragula*) is known in Australia as the Bar-tailed Goat-fish.

All these fishes are small, rarely exceeding 10 inches in length. Very little is known regarding their habits or distribution and they are not at present of much economic importance.

I have taken a few in a trawl near Penang and there is a possibility that with new methods of fishing they may become useful market fish.

Red Mullet are known to visit the British coasts, in vast shoals, at rare intervals.

BAT-FISHES.

(SCORPIDIDÆ.)

As far as I know, these fish are represented in our waters by the genus *Psettus* only.

The **Gēdabang** or **Nyior-nyior** (*P. argenteus*) is known in Australia as the Silvery Bat-fish. It attains a length of about eight or nine inches only and its breadth is about equal to its length. It is common and of fairly good edible quality but is not, at present, of importance.

The **Nyior-nyior** (*P. falciformis*) is also a small fish attaining a length of perhaps 9 or 10 inches.

CORAL FISHES.

(CHAETODONTIDÆ.)

A large group of about 200 species of marine carnivorous fishes, confined to the Tropics, mostly of small size and remarkable for their singular forms and markings and brilliant colours.

They are particularly abundant about volcanic rocks and coral reefs; but some ascend estuaries and tidal rivers, though not to any great distance.

The **Ketang** (*Ephippus argus*) ranges from the Indian Ocean to China and Australia, attaining a foot in length. If taken in the sea or in clean back-waters it is an excellent edible fish, but those captured in the vicinity of polluted rivers should be avoided, as there is evidence that they are foul feeders.

Hamilton Buchanan remarks of it, "When newly caught it is a fish of great beauty, easy digestion, and excellent flavour: but after death it soon becomes soft and strong tasting." In Ceylon "It is generally esteemed, its flesh partaking the flavour of trout" (Bennett).

This fish and its congener (*E. multifasciatus*) are favourably known in Australia as Butter-fish and are a common table fish in hotels and restaurants.

Ikan inggu or **Ikan babi** are Malay equivalents for the genus *Holacanthus*. The former term applying to the colouration and the latter to the rather pig-like profile and the presence, in all these fishes, of a pair of pre-opercular spines directed backwards, which are considered to resemble boar's tusks.

The **Bonang** (*Plataea teira*) is a deep-bodied fish which attains a length of at least 20 inches. Russell says their flavour is excellent and Cantor makes the same remark.

It is known in Australia as the Dark Bat-fish.

MOON-FISH.

(*DREPANIDAE.*)

The **Daun bēharu** (*Drepane punctata*) is a very common fish of fair edible value.

Considerable quantities of this fish have been taken in trawls both in India and Ceylon. Sir K. Gupta says they are very much sought after and always command a good price in Bengal. They are rather too bony to be popular with Europeans in the East.

“BLACK TREVALLY.”

(*TEUTHIDIDAE.*)

According to Dr. Boulenger's classification (1902), this family comprises a single recent genus, *Teuthis*, with about 30 species, herbivorous fishes from the Indian and Western Pacific Oceans. According to Bottard (“Poissons venimeux,” Paris 1889) the sting from the spines of these fishes is much dreaded, and this I can vouch for, though personally I have suffered very little inconvenience from the pricks of these spines.

It will be noticed that Duncker gives the generic synonym **Ketang** to members of this family and this is the name given by Malays to the genus *Ephippus* (*CHAETODONTIDAE*) which also has venomous spines.

In all species of *Teuthis* there are 13 dorsal spines and 7 anal spines, whereas *Ephippus* has 9 dorsal and 3 anal spines, which shews that the Malay system of classification does not agree with that adopted by scientists.

The **Dengkis** (*T. nebulosa*) is known on the East coast of Australia as the “Black Trevally” and the **Debam** (*T. java*) is also found on the Australian coast.

They are small fish, fairly common in the markets where they find a ready sale.

GOURAMI, ETC.

(*OSPHROMENIDAE.*)

This family of fresh water fishes is remarkable for several reasons:

From an edible point of view, because it includes the **Kalui** (*Osphromenus olfar*), known in India as the **Gourami**, which has a world wide reputation as one of the finest flavoured fresh water fish known, as well as the **Pēpuyu**, a favourite food fish in Negri Sembilan.

From an athletic and sporting point of view, because it includes the **Pêpuyu** (*Anabas scandens*) the famous climbing Perch, mentioned in all natural history books, as well as the **Ikan bēlaga** the equally famous fighting fish, on which Pahang rajahs have won and lost fabulous sums; and from a scientific point of view, because all members of this family are provided with super-branchial respiratory organs, situated in a cavity above the gills which enables them to live, happily, out of water for long periods.

The **Kalui** grows to a length of about two feet and is regarded as one of the best flavoured fishes in the East. It has been acclimatised in India, the Guianas, Mauritius and Australia.

Day writes,

“Commerson who observed it in the Mauritius in 1770, states that he never ate any fish more exquisite in flavour, whether from the sea or fresh water: he also added that in Batavia the Dutch reared them in large earthen pots, changing the water daily and feeding them on nothing but fresh water plants, especially the *Pistia natans*.”

General Hardwicke¹ gives an account of the breeding of this fish, apparently monogamous; he observes,

“They commence at six months of age, whilst their fecundity is astonishing. During the breeding season, they frequent the sides of tanks, where shelter is afforded them by the grasses and weeds growing in the water. For several days they are very active, passing in and out of their grassy cover, and in some places thickening it, by entangling all trailing shoots, and forming what is generally considered the spot under which the ova are deposited. They continue to watch this place with the greatest vigilance, driving away any interloping fish, and, at the end of a month numerous fry appear, over which the old gouramies keep watch many days.”

I kept these fish in a large pond at Kuala Pilah, having first caught them with a casting net in the Muar river. Their natural food consists of aquatic plants and I used to collect the leaves they like and send a leaf at a time down the stream until a **Kalui** rose to the bait. It was then a simple matter to lure the fish nearer and nearer, with carefully flicked leaves until it was close enough to my place of concealment to enable me to throw the net over the floating leaf under which the fish was rising.

There are probably many old friends who will remember the little dinners in Kuala Pilah, when the fish, fowl, mutton and vegetables were all locally raised.

The **Kalui** in my pond were fed daily on leaves, principally wild caladium and tapioca shoots, not thrown broad-cast but inserted in split bamboo poles which were pushed into the bottom of the pond. They ate a tremendous lot and grew very rapidly;

¹ Zool. Journ. IV, p. 309.

the caladium leaves imparting a very fine flavour to the fish.

They will rise to a fly or beetle, and some flowers, particularly a large Hibiscus. Anyone intending to keep these fish in stock ponds is advised to keep the pond free from pollution and to feed the fish regularly. It is only in this way that rapid growth and good flavour can be obtained.

They attain a length of two feet, a weight of at least 20 pounds and in shape resemble the turbot.

The **Pěpuyu** or **Bětok** (*Anabas scandens*) has a world wide reputation as the Climbing Perch. Gunther¹ tells us that in 1797 Daldorf in a memoir communicated to the Linnean Society of London mentions that he had himself taken, in 1791, an *Anabas* in the act of ascending a palm tree (*Palmyra*) which grew near a pond. The fish had reached the height of 5 feet and was going still higher. He goes on to say that the fish is named in the Malayan language the "Tree Climber," which is a mistake. He should, I think, have said the Malayalam language. See Day (Fishes of India) *Undi colli*.

Meek² writes,

"*Anabas* has been frequently obtained on the ground and a specimen now in the collection of Armstrong College, obtained from near Bangkok was found crossing the road 50 yards from the nearest water. It is named the climbing perch from the habit it has of climbing up the rough bark of trees by movements of the spine-clad opercula.

"The method of progression out of the water and the climbing of palms and palmyra trees, especially after heavy rains, have been repeatedly observed."

The Negri Sembilan Malays have a saying, often quoted, which hits off the high estimation in which this little fish is held by inland dwellers: *Jikalau sudah minum ayer gopong bertali ijok, sudah makan pepuyu, payah nak tinggalkan negri ini*: which may be roughly translated: When a visitor has drunk the water and eaten the fish of this country, he is loath to leave it.

The Ikan **Pělaga** or **Bělaga** (*Betta* spp.) probably derives its name from Siam where it is known as Pla Kat (*Pla*, fish; *Kat*, a fighter).

It is common throughout the Peninsula and may be caught in most of the ponds and ditches in Singapore.

Cantor relates that the Siamese are infatuated with the combats of these fish, staking on the issue considerable sums, and sometimes their own persons and families.

The licence to fight these fish used to be farmed in Siam and brought in a considerable revenue to the King.

¹ Study of Fishes, p. 516.

² Migrations of Fish, 1916.

The male fish are kept in bottles separately, and when in a state of quiet they are dull looking little fish, but if two bottles be brought together, the little creatures become greatly excited and the raised fins and whole body shine with bright metallic colours of dazzling beauty,

If two male fish are then placed together they fight like terriers. When fighting they utter a curious ringing note which sounds like "Kring Kring" and probably this accounts for another name by which they are known viz. **Ikan karing**.

The **Kēpar** (*Polyacanthus hasseltii*) is another beautiful little fish and quite common in brackish swamps and ponds.

This fish has been bred in confinement by Chinese, probably for centuries, and is known as the Paradise fish to aquarium owners in Europe. In its native element, in dark or muddy water, it is of rather a drab brown colour but if kept in a bowl in clear water, it has a beautiful golden colour with red transverse bands.

CORAL FISHES.

(POMACENTRIDAE.)

This family resembles the CHAETODONTIDAE (Coral Fishes) in form and mode of life, likewise in the brilliant colouration. For this reason I have applied the same English name in the absence of any other for this particular family.

Over 150 species are known. Some 30 species are described in Day's Fishes of India and probably the family is better represented in Malayan than in Indian waters.

As the names **Inggū** and **Gombing** shew, the Malays include CHAETODONTIDAE and POMACENTRIDAE in one family and curiously enough, the scaly-finned fishes (CHAETODONTIDAE) resemble the POMACENTRIDAE so closely that in some instances actually the same colouration and markings are common to members of the two families. This, as remarked by Dr. Günther, is one of many instances shewing that the colouration of animals depends to a great extent on their mode of life and natural surroundings.

From an edible point of view they are not of much economic importance but all specimens brought to the markets seem to find ready purchasers.

"WRASSES" OR PARROT FISHES.

(LABRIDAE.)

The "Wrasses" form a large family of most brilliantly coloured marine fishes, inhabiting all the tropical and temperate coasts.

Their regime is partially herbivorous, partially carnivorous. About 400 species are known.

Some of the members of this family have been observed to build nests for the protection of their eggs and young.

These nests in the European *Labrus* are made of sea-weeds, zoophytes, corals, broken shells, etc., and are the work of both the male and female. It is also in this family that sleep was first observed in fishes, and this has been fully verified by Mobius, on *Labrus rupestris* in an aquarium, the fish seeking a sleeping place at night and laying itself down to rest on one side.¹

Tokak is the generic name applied by Malays to those members of this family which are provided with strong canine-like teeth. (See Wilkinson's Dictionary, p. 201. **Tokak**. Biting, used of dogs, sharks, tigers, and other animals which use their teeth as a weapon of offence.)

The teeth of these fishes are used however for crushing shells, coral, etc.

A **Tokak** (*Chaerops omnopterus*) is known in New South Wales and Queensland as the Blue-spotted Groper.

Little use is made of this large family of fine edible fishes from a commercial point of view.

Their capture is confined to the hand line and to basket traps. Their habitat, deep water in the vicinity of coral reefs, renders the use of ordinary nets impossible but the trammel net which is unknown in this region should prove effective.

Many members of this family attain a weight of 50 pounds.

PARROT-WRASSES.

(SCARIDÆ.)

This family is closely allied to the preceding, with which they have been usually united, but differing in the more or less coalescent teeth, forming, often, a parrot-like beak.

I have placed the **Běchok** in this family and also among the LABRIDÆ as there are several species.

Mr. A. W. H. Hamilton, who is an authority on Malayan sea-fishes, tells me that the Malays of the Western part of Singapore confine the synonym **Běchok** to a fish with green teeth, which seems to identify his fish as *Pseudodax moluccanus* (Day, 2nd edition Vol. II, p. 421).

HORSE MACKERELS.

(CARANGIDÆ.)

A large and important family of carnivorous fishes, all of which are edible and many of large size.

Members of this family compose the bulk of the fish taken in nets on the East coast, which are dried and salted for export.

Some of our principal local varieties are the **Chěncharu** (*Caranx rottleri*), the **Sělar** (*Caranx*, not less than 12 species), the **Chermin** (*C. gallus*), the **Nyior nyior** (*Trachynotus* spp.) and the **Talang** (*Chorinemus* spp.).

¹ The Cambridge Natural History, 1904.

When freshly caught and cooked they are all excellent eating, but they do not keep well.

The **Chēncharu** is quite common and is found in large shoals. When in season, large numbers will be found in the markets and if quite fresh they are good edible fish. They are said to attain a length of 5 feet.

All the **Sēlar** are good eating, but the fresher they are the better. They give good sport with a light rod and small hook, to which a few small white feathers have been "whipped."

They like shade and will be found in the neighbourhood of piers and under vessels. When cruising, I have often noticed **Sēlar** taking shelter under my yacht, when we were becalmed, and if the period coincided with a meal time, we used to catch as many as we wanted in a few minutes.

There are, at times, large numbers of **Sēlar** in shallow water off Singapore as, probably, many sea-side residents know.

The **Chermin** (*C. gallus*) is a deep-bodied fish somewhat resembling the Dory in shape and is one of the best, if not the best, food fish in this family. It is found generally on reefs; takes a prawn or fish bait, and gives splendid sport as it fights very hard and takes a bit of playing.

Specimens 2 feet in length are not uncommon and it is said to grow to five feet in length. It is known in Australia as the Silvery Moon-fish.

The **Nyior nyior** (*Trachynotus ovatus* and *T. bailloni*) are known in Australia as the Dart. These fish must be fresh to be appreciated.

The **Talang** (*Chorinemus* spp.) is a common fish in the markets and fairly popular with most Asiatics, but some Malays have a prejudice against it and will tell you that it gives them an irritating and disfiguring affection of the skin.

It may be that the general appearance of these fish, all of which have a row of dark blotches along the side, may suggest the unsightly blotches seen on the faces, bodies and limbs of natives who are afflicted with certain kinds of skin disease, *kēdal*, *sopak*, etc. or that the consumption of this fish when not perfectly fresh causes urticaria, but the subject should be worth investigation.

One local species of **Talang** (*S. sancti-petri*) is known also in Australia as the Queen-fish, and another (*S. tooloo-parah*) is known also in Philippine waters as the **Talang-talang**.

There are many other excellent food fish in this family including the **Lēmbudok** or **Dēmudok**, **Gērēpoh** and **Berkas**, not specifically identified.

The generic term by which the *Caranx* branch of this family is known to the Moros, in the Philippines, is **Daing puti**. The name **Daing bēlang** occurs locally and is applied to *Caranx speciosus* and *C. compressus*,

MACKERELS, TUNNIES, ETC.

(SCOMBRIDAE.)

The fishes of the "Mackerel" family are pelagic forms, abundant in all the seas of the tropical and temperate zones. They travel about in shoals, spawn in the open sea, but periodically approach the shore in pursuit of other fishes on which they feed.

Our most important local members of this family are the **Pělata** (Siamese *pla thu*) (*Scomber microlepidotus*), the **Tongkol** (*Thynnus thunnina*) and the **Těnggiri** (*Cybium* spp.).

The **Pělata** is a fish of great commercial importance in Siam and on the East Coast, where it is extensively salted and dried for export.

The **Tongkol** is the Malay generic term for the Tunny and, I believe, for the Bonito also. These fish gives excellent sport when they are on the feed but often one sees a school of these fish jumping and disporting themselves, and on such occasions they seem to disdain the bait which is "trolled" past them.

The **Těnggiri** is, in my opinion, the best fish in our waters. The best both from a sporting and from an edible point of view, but I may be prejudiced in its favour because I have had more sport with this fish than with any other. Seale¹ gives corroborative evidence as to its edible qualities as follows, "In this family is the **tanguingue**, which is a true Spanish Mackerel. By many people this is regarded as the finest food fish in the Philippine waters."

A recent visitor to Singapore from Queensland told me that he had had great sport with these fish on the Barrier Reef and that they attained a weight of 100 pounds.

The big fish stay out in deep water and the best time to take them is during the N. E. monsoon. The best bait is a whole fish about 8 or 9 inches long, and at least 100 or 150 yards of line should be run off the reel, so as to keep the bait a long way astern as you sail along in a good breeze.

When making a passage in a heavy sea with no time for rod fishing we used to boom out as many as five brass wire lines and perhaps have two or three fish on at once averaging 20 pounds or so.

When our fisheries are better understood and depots with refrigerating plant are established on the islands off the East coast, more attention will certainly be paid to our oceanic fishes. Sea going fishing smacks should do a good trade with catches of Bonito, Tunny and Spanish mackerel.

One of the Spanish mackerels in America is one of the most highly esteemed of all American fishes and always commands a high price. Stead mentions that the catch in 1897 amounted to 1,183,456 pounds, worth nearly £14,000.

¹ Fishery resources of the Philippines.

HAIRTAILS.*(TRICHIURIDAE.)*

The **Timah-timah** (*Trichiurus* spp.) are some of our commonest fishes and are generally on sale in the markets.

I have never eaten them but the Chinese and Indians purchase them readily.

These fish have no caudal fin, the body being ribbon like and tapering to a fine point.

Miniature specimens an inch or two in length form a considerable proportion of the catches of illegal purse nets. The ordinary size of marketable specimens is about three to four feet.

Day quotes Russell as observing that in his time they were esteemed by the European soldiers in India, and Jerdon states that they afford very delicate eating.

SAIL-FISHES.*(HISTIOPHORIDAE.)*

A family of large oceanic fishes, occurring in tropical or sub-tropical seas. On account of their formidable sword, large specimens are held in dread by fishermen and are rarely taken and still more rarely preserved.

The Japanese in Hawaii have a regular fishery for Sail-Fish and Tuna. The Japanese fishermen in Singapore, who are the only deep water fishermen in our waters and whose methods are much more enterprising and thorough than those of the Malays and Chinese, are taking these fish occasionally.

I am informed that a Sail-Fish, three fathoms long was sold in the Clyde Terrace market within the past two weeks, but the information arrived too late to enable me to get a photograph.

This fish is known to Malays as **Sēlayer** or **Layeran** (*Layer*, a sail), and is by no means rare.

FLAT-FISHES.*(PLEURONECTIDAE.)*

Flat fishes are a large group of some 500 species, mostly marine.

The very young are transparent and symmetrical with an eye on each side, and swim in a vertical position like other fishes.

As they grow, the eye of one side moves by degrees to the other side, where it becomes the upper eye.

If, at that age, the dorsal fin does not extend to the frontal region, the migrating eye simply moves over the line of the profile; in other genera, the dorsal fin has already extended to the snout before the migration takes place, and the eye, passing between the

frontal bone and the tissues supporting the fin, appears to pass from side to side through the head, as was believed by some of the earlier observers.¹

As a food supply the flat-fishes are of great importance, the flesh of the majority being of excellent quality and flavour, and they are deservedly popular with Europeans in Malaya.

The family is represented in our waters by, certainly, not less than 19 species, of which 12 are included in the systematic list in this work.

The Malay generic terms are **Ikan sabēlah** and **Ikan lidah** for all members of this family, but in some districts the name **Sabēlah** is applied to those genera which have a distinct caudal fin (*Psettodes*, *Pseudorhombus*) and the name **Lidah** to the tongue-shaped genera (*Synaptura*, *Plagusia*, *Cynoglossus*).

In the Straits of Malacca these fish are very common in shallow water on sand and mud where they keep close to the bottom. This habit of keeping close to the bottom renders them particularly liable to capture by the beam or "Otter" trawl. I have taken these fish in a beam trawl in fair numbers both off Singapore and off the Krian coast.

On the great Kra flats off Krian which are formed of very soft mud I found it necessary to fit "ski" or wooden skates to the irons of the trawl to enable the trawl to slide on the surface of the mud, and took considerable numbers of these fish as well as some large Rays.

I should expect a trawler to be successful on the long banks and in the deep water gullies which, as a glance at the chart will shew, run in the direction of the prevailing currents, in many parts of the steamer route between Penang and Singapore.

An enormous amount of destruction of immature flat fish takes place daily in shallow water, specimens an inch or two in length being taken in seine nets and purse-nets from one end of the Straits to the other. A special effort should be made to stop this murder of miniature fish which has diminished our food supply to a very considerable extent.

Two species of our **Ikan sabēlah** are found on the Queensland coast. One, *Psettodes crumei*, is known as the Queensland Halibut, and the other, *Pseudorhombus russelli*, is generally called the "Flounder."

GOBIES.

(GOBIIDÆ.)

A large family of some 600 species, the great majority marine, mostly carnivorous and of small size.

The largest form (*Eleotris marmorata*) from the rivers of Siam, Borneo, Sumatra and the Malay Peninsula grows to nearly three feet, whilst the smallest (*Mystichthys luzonensis*) from the Philip-

¹ Cambridge Nat. History.

pinces, attains a length of about half an inch and is believed to be the smallest known fish.

The family is not of much economic importance at present and I have no personal knowledge of their edible qualities.

Our most noteworthy species are the **Běłontok** (*Eleotris marmorata*) the **Běłodok** (*Gobius* spp.), the **Těmbakul** and **Běłachak** (*Periophthalmus* spp.).

The **Běłontok** has already been alluded to as attaining a large size and not less than seven species are known to inhabit our waters.

One of our **Běłodok** (*G. butis*) is said by Day to be much esteemed by the natives of India, as being very light and wholesome, but unless elaborately cooked is not relished by Europeans, because of its deficiency in, or earthy, taste.

It attains a length of a foot and a half, takes a bait freely and is largely bred in tanks in India.

The **Těmbakul** and **Běłachak** will be familiar to most residents in Malaya as the Mud-Skippers which may be seen disporting themselves on the mud and among the mangroves, along all our coasts and estuaries.

Malays have told me that these fish are good eating and possess great medicinal virtues.

They have very conspicuous prominent eyes, which are capable of protrusion and retraction, and extraordinary muscular pectoral fins which they use like arms for progression on mud and for climbing.

Day writes, "They climb on to trees, holding on by their pectoral fins exactly as if they were arms. Now and then they plant these firmly as organs of support, the same as one places one's elbows on a table, then they raise their heads and take a deliberate survey of surrounding objects."

Saville-Kent is quoted by Stead as follows:

"A remarkable circumstance associated with the life economy of *Periophthalmus* is the fact that it cannot sustain life if continually water-submerged like ordinary fish. The exposure of its tissues to the action of atmospheric air with every fall of the tide appears to be essential to its well-being, and examples experimentally kept under water for prolonged intervals were literally drowned.

"As a provision for its abnormal life-habits, it has been ascertained that *Periophthalmus* possesses a supplementary respiratory organ which, singular to relate, is represented, in this instance, in the creature's tail.

"The fish while reposing on the surface of the mud commonly leaves its tail more or less immersed in the water. The blood circulates with abnormal energy through this thin membranous appendage, which accordingly fulfils the function of a supplementary gill."

SUCKING FISHES.

(ECHINEIDIDAE.)

These fishes, generally known as *Remora*, attach themselves by means of a remarkable adhesive disc on the upper surface of the head to boats and ships, or to whales, sharks and turtles and in this way manage to do a good deal of travelling with the minimum amount of effort. As they are not strong swimmers they obtain a much larger supply of food by riding about in this way than otherwise would be possible.

The natives of Cuba, Zanzibar and the Torres Straits use these fish for catching turtles; the fish being held by a metal ring round the base of the tail to which a line is attached. "When one of these fish, a foot in length, has its wet sucker applied to a table, and is allowed time to lay hold, it adheres so tightly that it is impossible to pull it off by a fair vertical strain" (Lydekker)¹.

The **Gēmi** (*Echineis naucrates*) is very common in these seas. It takes a bait readily, is edible, and may, occasionally, be seen in the markets.

GOBLIN-FISHES.

(SCORPAENIDAE.)

Some members of this family are Perch-shaped and edible, growing to a large size (*Sebastes*, *Scorpaena*, etc.).

Nearly all are distinguished by a powerful armature, either of the head, or fin spines, or both, and in some the spines are provided with poison glands (*Scorpaena*, *Pterois*, *Pelor* and *Synancia*) and a sting from these spines is extremely painful.

Lēpu is the Malay synonym for all members of this family.

FLAT-HEADS.

(PLATYCEPHALIDAE.)

This family with a single genus, *Platycephalus*, and some 40 species, inhabits the coasts of the Indian Ocean and the Western Pacific.

The Malay generic term is **Baji-baji**, so called from the wedge shaped head, and so far some four species have been identified in Malayan seas.

They live on the bottom, hidden in the sand as a rule, and as they depend on their protective colouring and spines to save them from possible enemies, they do not swim to any distance when disturbed but dart away for an instant and then lie motionless half buried in the sand.

This peculiarity renders them particularly liable to be taken by trawls and a large proportion of the catches made by the New South Wales trawlers is composed of these fish.

They are good edible fish and common in the markets.

¹ Royal Nat. History.

"STAR-GAZERS."

(LEPTOSCOPIIDAE.)

Information is wanting, but, I think the **Pukul gendang** (*Percis pulchella*) is rare and economically unimportant.

SPINY-EELS.

(MASTACEMBELIDAE.)

These are eel-shaped carnivorous fishes, very common throughout Malaya where they are known by the generic term **Tilan**. The largest species reach a length of three feet and the flesh of all species is of excellent quality. They are found far inland and often at considerable elevations.

Day states, "Excellent as food, although owing to their resemblance to eels (in fact they are eels with spines) or snakes, some people object to them." Buchanan observes, "sought after by the natives, the highest of whom in Bengal make no scruple of eating them; and by Europeans they are esteemed the best of the eel-kind."

FROG-FISHES.

(BATRACHIDAE.)

These carnivorous fishes apparently delight in mud and dirty water; they frequent the shores, ascending tidal rivers and estuaries. At Penang "the natives attribute poisonous qualities to these fishes, and reject them even as manure" (Cantor).

ANGLER-FISHES AND "CROAKERS."

(ANTENNARIIDAE & MALTHIDAE.)

These fishes have no economic value.

LEATHER-JACKETS.

(TRICANTHIDAE AND BALISTIDAE.)

These two families may be conveniently taken together in this small work as there is a strong affinity between them.

Though containing many species of no economic value one species, the **Jëbong** (*Balistes stellatus*), is preferred to all other fish by many Malays, including fishermen, whom I have questioned. I think that the main reason for this preference is that the flesh of this fish more nearly resembles that of a chicken than any other fish, and consequently the change to what approximates to a meat diet is welcomed.

The **Jëbong** has a tough leathery skin which has to be removed before it is cooked. The cook should not be allowed to remove the head which is the best part of this fish, and of many others, especially perhaps the **Tenggiri**.

Leather-jackets are held in considerable esteem in Australia as food fish and are commonly served in hotels and restaurants.

BOX-FISHES.

(OSTRACIONTIDAE.)

This family is of no edible importance.

GLOBE-FISHES AND PORCUPINE FISHES.

(TETRODONTIDAE & DIODONTIDAE.)

These fish possess poisonous properties and instances have been recorded of persons dying shortly after eating them. Malay fishermen, however, commonly eat the **Buntal pisang** (*Tetrodon lunaris*) and some other species, being careful to remove all the poisonous organs.

SHARKS AND DOG-FISHES.

(CARCHARIIDAE, SCYLLIDAE, SPHYRNIDAE.)

Sharks are active predacious fishes living at different depths in the sea from the surface to nearly a thousand fathoms and ranging from mid-ocean to the shallower waters round the coasts in every part of the world. They are most abundant in the Tropics where they attain their greatest size, and some of the Sharks are the largest of living fishes.

Among the SCYLLIDAE (Dog-fishes) we have in these waters the Tiger or Zebra Shark (**Yu chechak** or **Yu to'kek**) with dark bands on a tawny ground which attains a length of at least 10 feet.

Among our species of the true Sharks (CARCHARIIDAE) we have **Yu tenggiri** (*Galeocerdo rayneri*) which attains a length of over 12 feet and is very ferocious, but fortunately rather rare, and the **Yu jerong** or **Yu sambaran** (*Carcharias* sp.) which has also a bad reputation.

The Hammer-head Sharks (SPHYRNIDAE) **Yu bengkong**, **Yu sanggul** or **Yu palang** are voracious, usually live in deep water and grow to a length of 15 feet.

There is no scientific record of the appearance of RHINODONTIDAE in these waters. Sharks of this family are probably the largest known and are said to exceed 50 feet in length (some writers mention 70 feet), but to be quite harmless. Specimens have been seen or captured in the neighbourhood of Ceylon, and on one occasion I watched a very large shark, in clear water, near Nipah Bay, Tioman Island, for more than half an hour, which appeared to equal the length of my yacht (35 feet).

The economic value of sharks has not yet been fully realised. Fishermen regard them as a nuisance as they tear nets and take fish off their hooks, and they are avoided as much as possible. Incidentally sharks are a nuisance to trawl-fishermen in Australia, and if there is any delay in getting the "cod-end" containing the fish on board, the sharks will bite pieces out of it. There is however every indication that shark-leather will soon be an ordinary trade commodity. The skin of sharks is composed of two layers:

the outermost integument, "shagreen," is covered with denticles, and hitherto, owing to the difficulty of treatment, has had a very limited use, but within the last few years a method has been discovered of separating the outer and inner skins and the latter can be tanned and used in every way like ordinary leather. It is therefore likely that the high price and scarcity of ordinary leather will eventually lead to the universal exploitation of the shark, ray and porpoise fisheries with special nets and appliances. I see in the Australian Magazine "Sea, Land and Air" (September 1920) that a Marine-Leather Company is operating successfully off the coast of Florida and North Carolina.

Other commercial products are the blood, fins, liver and meat. The blood is said to furnish one of the finest waterproof glues yet known for aeroplane propellers, etc.; the fins are a well known Chinese delicacy, and the American Bureau of Fisheries has published some thirty recipes for cooking shark-meat.

Small sharks are esteemed as food by the Malays, Indians and Chinese and are excellent eating.

The liver of the shark is rich in oil and is said to equal that of the Cod in its medicinal properties. It is also used in the preparation of soap, paint, etc., including the treatment of leather.

SAW FISHES.

(*PRISTIDAE*.)

The family contains one genus (*Pristis*) with about four or five species.

These fish are termed **Běroi** by Malays in some districts but the descriptive names **Yu gergaji**, **Yu parang** and **Yu todak** are more commonly heard, Malays placing these and the RHINOBATIDAE among the Sharks (SELACHOIDEI) and not among the Rays (BATHOIDEI), with good reason.

Boulenger states that an arbitrary distinction has been made which has little to recommend it except custom and some measure of convenience.

These fish are readily eaten by Malays, Chinese and Tamils and are very common. They enter rivers right up into fresh water and small specimens two or three feet long are often taken accidentally in casting nets.

They have always appeared to me to be very lethargic and sluggish and as the small ones in a net give less trouble than any other fish of the same size, I have always considered them to be more formidable in appearance than in reality. However, Day writes "Great injuries are inflicted by these fishes, which strike sideways with their formidable snouts; and although not personally a witness to the fact, I have been informed on native authority, that large ones have been known to cut a bather entirely in two."

It would be interesting to know whether there is any record of patients having been admitted to hospital in India or Malaya, suffering from injuries inflicted by these fish.

A saw-fish measuring 23 feet 6 inches exclusive of the saw was taken in the Bay of Bengal by the Government trawler "Golden Crown" and I believe that this is the largest recorded fish. No mention is made of the length of the saw of this specimen but it is not likely to have been less than 7 feet. The largest saw in the Raffles Museum, Singapore, measures 5 ft. 10 $\frac{3}{4}$ in.

BEAKED-RAYS.

(*RHINOBATIDAE.*)

These are harmless, sedentary, bottom-feeding fishes which subsist chiefly on shell-fish, crabs, etc. They are considered good eating and are sold regularly in the markets.

They are known to Malays as **Yu kēmējan**.

ELECTRIC-RAYS.

(*TORPEDINIDAE.*)

These Rays to which the Malays have given the descriptive names **Pari kēbas** or **Pari sēbar** have the power of inflicting electric shocks. "The fish" writes Dr. Günther, "gives the electric shock voluntarily, when it is excited to do so in self defence, or intends to stun or kill its prey. The electric currents created in these fishes exercise all the other known properties of electricity; they render the needle magnetic, decompose chemical compounds, and emit the spark."

Our Malayan species are very small. I have a specimen of the **Pari kēbas** (*Astrape dipterygia*) about six inches long and there is no record yet of specimens over 18 inches.

When trawling on the Australian coast we took many specimens which appeared to be between two and three feet in length and one or two new deck hands experienced shocks which appeared to cause only momentary inconvenience.

Cantor says that out of the water they may be handled with impunity.

STING RAYS.

(*TRYGONIDAE.*)

Nearly all the members of this family are provided with long whip-like tails, which are generally armed with spines. In the larger kinds these formidable spines may be as much as 8 or 9 inches in length; and, as they wear out they are, from time to time, shed and replaced by new ones growing from behind.

These spines inflict very severe wounds, the pain of which is greatly increased by the apparently poisonous cutaneous mucus introduced into the wound.

The **Pari bēting** (*Trygon uarnak*) attains a large size, 5 feet or more across the disk, and a weight of well over 200 pounds. In one haul of the trawl in the Bay of Bengal the "Golden Crown" took four of these fish which weighed respectively 180, 170, 160 and 122 pounds.

The **Pari dēdap** (*Urogymnus asperimus*) is the sole representative of a genus and remarkable from the fact that its back is covered with osseous tubercles, among which are studded, at intervals, a number of conical denticles or spines rather like limpets in appearance.

This fish ranges between the Red Sea, East Coast of Africa, seas of India and the Malay Archipelago.

I recently overheard a Malay in the Raffles Museum apply the name **Dērēdap** to this fish and perhaps a note on the word **Dēdap** and its derivatives may be of interest.

Dēdap—a tree (*Erythrinus* sp.) with scarlet flowers, the bark of which is studded with spines of the same limpet-like shape as those of the **Pari dēdap**.

Dēdap—a shield or buckler.

Rēdap—a small drum, (probably so called from the kind of skin used).

Mērēdap—(Riau, Johor) springing up plentifully, of prickly heat and other skin eruptions, the feature of which is a large number of pustules.

The word **dēdap** as meaning a shield is obsolete both in colloquial Malay and in literature and it is interesting to note that its place has been taken by the Indian word *Pērisai*.

EAGLE-RAYS.

(MYLIOBATIDÆ.)

This family contains five genera and about 27 species. All five genera are represented in Malayan seas.

These fish feed principally on Molluscs, the shells of which they grind with their large grinding-teeth. Some of them attain an enormous size, over 20 feet in width, a thickness of 3 to 4 feet and a weight, probably, of over a ton.

They are variously known as Devil-fishes, Sea-devils, Bat-fishes, Eagle-rays, etc., and it is interesting to note that the terms Bat and Eagle are taken from the Malay, viz. **Pari kēlawar** and **Pari lang**.

The largest of this family are the **Pari paus** (*Dicerobatis* spp. and *Ceratoptera* spp.).

I have seen these fish leap out of the sea to a height of perhaps 7 or 8 feet, time after time, coming down each time with a tremendous splash, and Malays have told me that the fish does this to shake off the *remora* which hang on to them in large numbers.

In conclusion I may add, that all the Rays and Skates are eaten by natives of the East, while the "wings" or fins are highly esteemed by the Chinese. Fishes of this order would form a considerable proportion of the catches of a trawler and would provide a cheap and valuable food, for which there is a constant demand, either fresh or salted.

Malayan Fishes.

PART II.

ALPHABETICAL LIST OF MALAYAN FISHES.

NOTE:—The letters and abbreviations inserted in brackets after the Malay name of each fish, refer to authorities for both the Malay and scientific synonyms.

Where no authority is given the writer accepts responsibility for the identity of those species.

LIST OF ABBREVIATIONS.

C. = Cantor. D. = Dennys. R. = Rowell. R. M. = Raffles Museum. Dun. = Duncker. Blkr. = Bleeker. M. W. and de B. = Max Weber and de Beaufort. S. = Sauvage. Wilk. = Wilkinson. C. and S. = Clifford and Swettenham. S. M. = Selangor Museum. Blgr. = Boulenger.

Alu-alu. *Sphyraena novae-hollandiae*.

„ *obtusata*.

„ *jello*.

Barracudas. Fam. SPHYRAENIDAE.

Members of the genus *Sphyraena* are called “Barracudas” in America and elsewhere, and Pikes in Australia.

See also **Kachang-kachang** and **Těnok**.

Ambu-ambu (Wilk.). “The name of a large deep-sea fish. When preserved this fish is known as **Ikan kembal mas**.”

Probably **Tongkol** or **Ikan ayer**.

Thynnus thunnina or the Bonito *Thynnus pelamys*.

Mackerel. Fam. SCOMBRIDAE.

See **Kembal mas**.

Ampas tēbu (R. M.). *Pristipoma operculare*.

“Grunters.” Fam. PRISTIPOMATIDAE.

Ampit. **Anak ampit** (Wilk.): (Kedah) a fish; better known as **Ikan pēlaga**. This is probably the well known fighting fish **Ikan bēlaga**: *Betta pugnax*.
„ *bellica*.

Fam. OSPHROMENIDAE.

See also **Pala** and **Bélagá**.

Anding. See **Bélanak**. Grey Mullet. Fam. MUGILIDAE.

Anjang-anjang (Blkr. **Andjong-andjong**).

Pentapus caninus.

(R. M.) *Scolopsis ghanam*.

Grunters. Fam. PRISTIPOMATIDAE.

Aruan (Dun.). *Ophiocephalus guichua*

” ” *lucius*.

(Dun. C. D.). ” *striatus*.

The Murrel of Northern India.

The “Murrel.” Fam. OPHIOCEPHALIDAE.

Aruan tasek (Dun. D. R.). *Elacate nigra*.

Mackerel. Fam. SCOMBRIDAE.

Asoh-asoh (R.). *Lethrinus nebulosus*.

Sea-Bream. Fam. SPARIDAE.

Ayam (Blkr. **Hajam**). *Monacanthus choerocephalus*.

” ” *hajam*.

Balistes stellatus.

Leather-jackets. Fam. BALISTIDAE.

Ayer. **Ayer-ayer** (Cliff.). “The name of a sea-fish” *Thynnus thunnina* C. V. The **Tunny** or **Tuna**.

The name “**Ayer**” is used on the East coast of the Malay Peninsula, and **Tongkol** elsewhere.

Mackerel. Fam. SCOMBRIDAE.

Babi. *Holacanthus* spp.

So called from the shape of the head and the presence of a spine considered to resemble a pig’s tusk.

Coral-Fishes. Fam. CHAETODONTIDAE.

Bagat. *Caranx* sp.

Horse-Mackerel. Fam. CARANGIDAE.

Bagok. *Arius* sp. Cat-fish. Fam. SILURIDAE.

Baji-baji (D. R.). *Platycephalus tuberculatus*.

” *macracanthus*.

” (R. M.). ” *punctatus*.

Flat-heads. Fam. PLATYCEPHALIDAE.

Bakap (Unid.). Cat-fish family. SILURIDAE.

Bakok or **Bangkok**. q.v.

Balut (M. W. & de B. III 515). *Macrotema caligans*.

An eel belonging to the order SYNBRANCHOIDEA.

Bambangan. Also **Bambang** and **Mambang**. *Lutianus* sp.

Snapper. Sub-fam. LUTIANINAE.

Banang. Large of its kind: viz. **Puput banang**. **Jolong-jolong banang**.

Bandan (D. R.). *Sparus hasta*.

Sea-Bream. Fam. SPARIDAE.

Bandang (M. W. & de B. II 15). *Chanos-chanos*.

The Salmon-Herring of Australia.

The Milk-fish of India.

The Milk-fish. Fam. CHANIDAE.

(M. W. & B. II 3). *Elops hawaiiensis*.

The Bony-fish: A small relative of the Tarpon.

Giant-herring. Fam. ELOPSIDAE.

Bangkok. Also **Bakok** on East Coast.

„ (M. W. & de B. II 40). *Engraulis setirostris*.

„ „ *grayi*.

Herring. Fam. CLUPEIDAE.

Barat-barat (Blkr.). *Triacanthus strigilifer*.

„ *blockii*.

„ *nieuhofti*.

Monacanthus chinensis.

„ *penicilligerus*.

Leather-jackets. Fam. TRIACANTHIDAE.

Barau-barau. Also **Bēbarau** and **Sēbarau**.

Hampala macrolepidota.

(Dun. *Barbus hampal*).

Carp. Fam. CYPRINIDAE.

This fish gives good sport with live bait or spinner.

Barau-barau laut (R. M.). *Priacanthus hamrur*.

Sea Perch. Sub-fam. PRIACANTHINAE.

Batu (R. M.) (D.R.). *Proteracanthus sarissophorus*.

Sea-Bream. Fam. SPARIDAE.

Baung. The following varieties are recognised: **Baung akar**,

B. kunyet, **B. gantang**, **B. pisang**, **B. puntong**.

„ (Dun.). *Macrones nigriceps*.

„ (M. W. & de B. II 341). „ *nemurus*.

„ **kuning** (M. W. & de B. II 343). „ *planiceps*.

Cat fish. Fam. SILURIDAE.

Bawal (Tamil **Voval**). The following varieties are distinguished.

B. chermin (C.). *Stromateus atous*.

B. itam (D.). „ *cinereus*.

B. puteh. „ „

B. tambak (C.). „ *niger*.

B. kēdewas. „ *cinereus*.

Pomfret. Fam. STROMATEIDAE.

Bayan. Bayan-bayan.

(**Boyan** R. M.).

Chilinus fasciatus.

„ *chlorurus*.

Parrot-fish. Fam. LABRIDAE.

Běbēras (M. W. & de B. III 1g6). *Cyclocheilichthys apogon*.

Carp. Fam. CYPRINIDAE.

Běchok (Wilk. 93). *Julis lunaris*.

Parrot-fish. Fam. LABRIDAE.

Pseudoscarus spp.

Pseudodax moluccanus.

“Parrot-wrasses.” Fam. SCARIDAE.

Běgahak (J. S. A. S. VIII 120). *Belodontichthys dinema*.

Cat-fish. Fam. SILURIDAE.

Běkil. Also **Berkil**, q.v.

Bělachak. *Periophthalmus koelreuteri*.

Goby. Fam. GOBIIDAE.

Bělaga. Or **Berlaga**. **Ikan berlaga** (literally) fighting fish.

Betta pugnax.

„ *bellica*.

Three varieties of fighting fish are recognised, viz.,
Anak karing, **A. sempila**, **Karing gajah** and they fight
only with members of the same species.

See also Bleeker's Atlas Vol. IX, *Betta picta*.

Osphromenus striatus.

Fam. OSPHROMENIDAE.

Bělalang. *Exocoetus oligolepis*.

„ *neglectus*.

„ *nigripinnis*.

„ *speculiger*.

Flying-fish. Fam. SOMBRESOCIDAE.

- Bělanak** (Dun. D. R.). *Mugil cunnesius*.
 (Dun.). „ *planiceps*.
 „ „ *speigleri*.
 „ „ *waigiensis*.
B. jěmpul (C.). „ *planiceps*.
B. rapang (R. M.). „ *bleekeri*.
B. anding. „ *borneensis*.
B. tamok. „ *waigiensis*.

Other Malay varieties are **Bělanak angin**, **B. pakau**,
B kěděra, **B. puteh**, **Puntong damar** or **Puting damar**,
 and **Pělong**.

Grey Mullet. Fam. MUGILIDAE.

Bělau. See **Sělangat bělau**.

Bělibas. See **Gělibas**.

- Bělida** (R. M.). *Notopterus notopterus*.
 „ (Dun.). „ *chitala*.

Feather-backs. Fam. NOTOPTERIDAE.

Bělidang or **Běledang**. (C. & S. Dict. 259) a salt water fish
 shaped like an eel (unid).

- Bělin** (R. M.). *Muraena (Gymnothorax) undulata*.
Pisoodonophis cancrivorus.
 Eels. Fam. OPHICHTHYIDAE.

Běliak mata (M. W. & de B. II 68 **Mata belo**).
Clupea (Alosa) kanagurta.

Běliak mata jantan. *Clupea (Harengula) moluccensis*.

Běliak mata kapak. *Pellona dussumieri*.
 Herring. Fam. CLUPEIDAE.

- Bělodok** (Dun.). *Apocryptes lanceolatus*.
 „ *Periophthalmus schlosseri*.
 „ *Gobius giuris*.
 „ *Boleophthalmus boddarti*.

Bělodok kěrapu (Dun.). *Gobius sadanundio*.
 „ **lobang** „ *Gobius sp. aff. caninus*.
 Goby. Fam. GOBIDAE.

Bělodok karang (D. R.). *PlatyGLOSSUS dussumieri*.
 Parrot-fish. Fam. LABRIDAE.

- Bělontok** (Dun.). *Eleotris butis*.
 (Wilk.). *Gobius viridipunctatus*.
 Goby. Fam. GOBIIDAE.
- Bělungkor.** *Saurida tumbil*.
 Cf. M. W. & de B. II 142 **Belanka** (Bintang).
 The Queensland Smelt.
 Fam. SCOPELIDAE.
- Bělukang.** *Arius leiotetocephalus*.
 (Dun. *Arius liocephalus*).
 Cat-fish. Fam. SILURIDAE.
 See **Pědukang**.
- Bělut.** *Monopterus albus*.
 (M. W. & de B. III 414). An eel belonging to the
 order SYNBRANCHOIDEA.
- Běngkalis.** Also **Měngkalis**.
Ikan běngkalis is another name for the **Ikan těr-
 bok**.
- Běngkongkong.** Also **Běkukong**, **Běkuku** and **Kuku**.
 (Blkr. **Běkukung** VIII 108). *Sparus hasta*.
 Sea-Bream. Fam. SPARIDAE.
- Běntulu** (M. W. & de B. III 209). *Barbichthys laevis*.
 Carp. Fam. CYPRINIDAE.
- Běras-běras.** *Kyphosus* spp.
 Drummers. Fam. KYPHOSIDAE.
- Běrchat** (**S. bertchat**). *Ophiocephalus gachua*.
 "Murrel." Fam. OPHIOCEPHALIDAE.
 See **Aruan**.
- Běrkas.** *Caranx* sp.
 Horse-mackerel. Fam. CARANGIDAE.
- Běrkil.** A dark red fish of the **Ikan merah** family, which fre-
 quents timber rather than reefs, *i.e.* near piles,
 piers, sunken barges, etc.
Lutianus sp.
 Snapper. Sub-fam. LUTIANINAE.
- Běroi.** Also **Yu gergaji**.
Pristis spp.
 Saw-fish. Fam. PRISTIDAE.

- Běsikor.** Also **Měsikor.** *Diagramma* spp.
Grunters. Fam. PRISTIPOMATIDAE.
- Bětok** (Dun.). *Anabas scandens*.
The well known climbing perch of natural history books.
Fam. OSPHROMENIDAE.
- Bětulu.** See **Běntulu.**
- Biang-biang** also **Měmbiang** (M. W. & de B. II 29). *Setipinna breviceps*.
Herring. Fam. CLUPEIDAE.
- Biji durian** (Dun.). *Osphromenus malayanus*.
Fam. OSPHROMENIDAE.
- Biji nangka** (D. R.). *Upeneus tragula*.
Red Mullet. Fam. MULLIDAE.
- Bilis** (M. W. & de B. II 16). *Stolephorus commersonii*.
" *tri*.
("White-bait") Herring. Fam. CLUPEIDAE.
- Bonang.** *Platax teira*.
Coral-fish. Fam. CHAETODONTIDAE.
- Bongkar karang.** Literally the reef lifter. A name applied to large members of the Ray family.
- Boyan.** See **Bayan.**
- Bujok.** *Ophiocephalus* sp.
"Murrel." Fam. OPHIOCEPHALIDAE.
- Bulan** or **Bulan bulan** (M. W. & de B. II 6). *Megalops cyprioides*.
Giant-Herring. Fam. ELOPSIDAE.
- Bulu ayam.** *Coilia dussumieri*.
" *quadrifilis*.
(Anchovy) Herring. Fam. CLUPEIDAE.
- Bulus bulus** or **Běbulus** (D. R. **Bolas-bolas**). *Sillago sihama*
" *maculata*.
The Whiting of Australia. Fam. SILLAGINIDAE.
- Bunga ayer** (C.). *Stolephorus indicus*.
Engraulis Russellii.
See **Bilis**.
("White-bait") Herring. Fam. CLUPEIDAE.
Note: **Bunga ayer**, are probably the larvae or young of valuable food-fishes in the *Leptocephalus* stage.

Buntal. A name applied to a large number of fishes belonging to the families OSTRACIONTIDÆ (Box-fishes) ; TETRODONTIDÆ (Globe-fishes) ; and DIODONTIDÆ (Sea-porcupines).

Buntal batu (R. M.). *Ostracion cubicus*.

„ **kotak** or **pēti**. „ *nasus*.

„ „ „ *cornutus*.

Buntal pisang. *Tetrodon lunaris*.

„ **duri** (Dun.). „ *reticularis*.

„ **landak.** *Diodon novemmaculatus*.

„ „ (R.M.). „ *hystrix*.

Chabok. See **Parang-parang**.

Chandong. *Opisthopterus tartoor*.

Raconda russelliana.

Herring. Fam. CLUPEIDÆ.

Chělek mata (D. R. **Chileh mata**). *Pristipoma maculatum*.

“Grunters.” Fam. PRISTIPOMATIDÆ.

Chěmpěras also **Těmpěras** (R. M. **Temporas**).

Cyclocheilichthys apogon.

Carp. Fam. CYPRINIDÆ.

Chěncharu also **Jaru-jaru** (Dun.). *Caranx rottleri*.

Horse-mackerel. Fam. CARANGIDÆ.

Chěnchodak. See **Todak**.

Chěrechek (Cliff. 351). A fresh water fish with bright scales and red fins.

(Cf. M. W. & de B. III 62 **Tjettjereh**).

Rasbora argyrotaenia.

Carp. Fam. CYPRINIDÆ.

Chěrmin. *Caranx gallus*.

The Silvery Moon fish of Australia.

Horse Mackerel. Fam. CARANGIDÆ.

Chěrmin. See **Bawal chermin**.

Daing bělang. *Caranx compressus*.

„ *speciosus*.

Horse Mackerel. Fam. CARANGIDÆ.

Darok-darok. (See C. & S. dict. 395).

Carp. Fam. CYPRINIDÆ.

Daun (S. M.). *Barbus oatesii*.

Carp. Fam. CYPRINIDAE.

Daun (Dun.). *Platax teira*.

Coral-fish. Fam. CHAETODONTIDAE.

Daun baharu (D. R.). *Drepane punctata*.

The Moon-fish of Queensland.

„ „ *Ephippus orbis*.

Coral-fish. Fam. CHAETODONTIDAE.

Debam. *Teuthis java*.

“Black Trevally.” Fam. TEUTHIDIDAE.

Dĕlah. *Caesio lunaris*.

„ (R. M.). „ *kuning*.

„ „ „ *pinjalu*.

Dĕlah karang (D.). *Caesio chrysozona*.

Sea-Bream. Fam. SPARIDAE.

Dĕmbudok. *Caranx* sp.

Horse Mackerel. Fam. CARANGIDAE.

Dengkis. *Teuthis nebulosa*.

„ (R. Dukas). *Teuthis virgata*.

“Black Trevally.” Fam. TEUTHIDIDAE.

Duri (Dun.). *Macrones nemurus*.

„ (D. R.). *Arius sagor*.

Cat-fish. Fam. SILURIDAE.

Engor-engor. *Macrones nemurus*.

Cat-fish. Fam. SILURIDAE.

Gabus (Wilk. 557). *Ophiocephalus* sp.

“Murrel.” Fam. OPHIOCEPHALIDAE.

Garing (M. W. & de B. III 152). *Labeobarbus tambra*.

Carp. Fam. CYPRINIDAE.

Gĕdabang (D. R.). *Psettus argenteus*.

The Silvery Bat-fish of Australia.

Bat-fish. Fam. SCORPIDIDAE.

Gĕlam (D.). *Psammoderma vaigiensis*.

Sea-Perch. Fam. SERRANIDAE.

• **Gělama** (C.). *Umbrina russellii*.

„ **panjang** (D.). *Otolithus argenteus*.

Sciaena spp.

The following varieties are distinguished: viz.:

G. panjang.

„ **papan.**

„ **China.**

„ **sêkang.**

„ **rapang.**

„ **batu.**

„ **itam.**

„ **perak.**

„ **batu keling.**

„ **lanjut.**

„ **kuning dada.**

„ **dahi tinggi.**

„ **chêrua.**

„ **pisang.**

Jew-fish. Fam. SCIAENIDAE.

• **Gělibas** also **Bělibas** & **Libas** (R. M. Bělibas). *Teuthis oramin*.

“Black Trevally.” Fam. TEUTHIDIDAE.

• **Gemang.** The synonym for large **Ikan sěmbilang**.

Plotosus spp.

Cat-fish. Fam. SILURIDAE.

• **Gěmang darat** (Dun.). *Silurichthys phaiosoma*.

Cat-fish. Fam. SILURIDAE.

• **Gěmi** also **Gědemi** and **Kěmi** (C.). *Echineis naucrates*.

Sucking-fish. Fam. ECHINEIDIDAE.

• **Gěrěpoh.** Like the **Sagai** but with thicker lips.

Caranx sp.

Horse Mackerel. Fam. CARANGIDAE.

• **Gěrětak lantei** (R. M. **Kertah lantei**). *Synagris japonicus*.

„ „ *Scolopsis personatus*.

„ „ (D. R.). *Lethrinus nebulosus*.

„ „ Sea-Bream. Fam. SPARIDAE.

• **Gergaji.** **Yu gěrgaji.**

(Dun. C.). *Pristis cuspidatus*.

Saw-fish. Fam. PRISTIDAE.

Gërut-gërut (Dun.). *Mesoprion* sp.

Snapper. Fam. LUTIANIDAE.

„ „ (D. Blkr.). *Pristipoma hasta*.

„ „ (Blkr. **Krot-krot**) „ *maculatum*.

„ „ „ **Krot**) „ *guoraca*.

“Grunteræ.” Fam. PRISTIPOMATIDAE.

Gombing (Cf. R. **Rombin karang**). *Heniochus macrolepidotus*.

Coral-fish. Fam. CHAETODONTIDAE.

„ *Glyphidodon coelestinus*.

Coral-fish. Fam. POMACENTRIDAE.

Haruan. See **Aruan**.

Hayam. See **Ayam**.

Inggü. *Dascyllus* sp.

„ (R. M.). *Pomacentrus albofasciatus*.

„ (D. R. **Ingu**). *Amphiprion ephippium*.

„ (R. M.). *Amphiprion frenatus*.

Coral-fish. Fam. POMACENTRIDAE.

„ (D. R. **Ingu rombin**). *Holacanthus sexstriatus*.

„ (D. R. **Ingu rombin**). *Holacanthus mesoleucus*.

Coral-fish. Fam. CHAETODONTIDAE.

Jahan. *Arius thalassinus*.

Cat-fish. Fam. SILURIDAE.

Jalu jalu (R. M.). *Caranx boops*.

Horse-mackerel. Fam. CARANGIDAE.

„ „ *Caesio pinjalu*.

Also known as **Ikan merah china**.

Sea-Bream. Fam. SPARIDAE.

Jampong (R. M.). *Chilinus chlorurus*.

Parrot-fish. Fam. LABRIDAE.

Jangas = Bandang.

Fam. CHANIDAE.

Jarang gigi (C.). *Otolithus maculatus*.

„ „ *argenteus*.

„ „ *ruber*.

„ *Collichthys biaurita*.

Jew-fish. Fam. SCIAENIDAE.

Jaru-jaru. See **Chëncharu**.

Jěboh. See **Tamban jěboh.**

Jěbong. (D. R.). *Balistes stellatus*.

“Leather jackets.” Fam. BALISTIDAE.

Jěmbědi. *Engraulis* sp.

Herring. Fam. CLUPEIDAE.

Jěmpul. See **Bělanak.**

Jěněhak. *Lutianus roseus*.

(Blkr.), „ *johnii*.

„ „ *sebae*.

„ „ *fulviflamma*.

„ „ *lioglosus*.

„ „ *argentimaculatus*.

NOTE:—The generic names **Ikan merah** and **Ikan jěněhak** are synonymous. The latter name being used in the north, (Penang and Kedah) and the former in the south, Singapore, etc.

Snapper. Sub-fam. LUTIANINAE.

Jěngkua (unid.).

Carp. Fam. CYPRINIDAE.

Jěnjalu see **Jalu jalu.**

Jěrong. See **Yu jěrong.** Shark. Fam. CARCHARIIDAE.

Jolong jolong or **Julong:** also **Jěnjulong** (Dun.). *Hemirhamphus cantoris*.

(Dun.). *Hemirhamphus buffonis*.

„ „ *limbatus*.

„ „ *pogonognathus*.

„ „ *fluvialis*.

„ „ (Blkr.). „ *far*.

Jolong-jolong banang. *Hemirhamphus far*.

Gar-fish. Fam. SCOMBRESOCIDAE.

Juara (Wilk. 235). An edible fresh-water fish.

(Cf. M. W. & de B. II 258 **juaro**). *Pangasius polyuranodon*.

Cat-fish. Fam. SILURIDAE.

Kachang-kachang. A fish similar to, but smaller than, the **Alu-alu**, q.v.

“Barracudas.” Fam. SPHYRAENIDAE.

Kachi. *Diagramma* spp.

“Grunters.” Fam. PRISTIPOMATIDAE.

Novacula spp.

Parrot-fishes. Fam. LABRIDAE.

Kakap also **Siakap**. The “Cock-up” of Europeans in India, whence the name by which this fish is known in Queensland was probably derived.

(Blkr.). *Lates calcarifer*.

Sea-Perch. Fam. SERRANIDAE.

Kalat (R. M.). *Pseudoscarus rivulatus*.

“Parrot-Wrasses.” Fam. SCARIDAE.

Kalui (Dun, D. R.). *Osphromenus olfax*.

Habitat—China and the fresh waters of the Malay Archipelago.

Naturalised in Mauritius, Cayenne, Australia and introduced into some parts of India, viz., near Calcutta, Madras and the Neilgherries. Attains 20 lbs. or more in weight and is excellent eating when kept in clean water. Known as Gurami in India.

NOTE:—**Kalui** probably derived from **Kallawah**. (Tamil) a perch.

Fam. OSPHROMENIDAE.

Kapas, Kapas-kapas, (Blkr.). *Sparus hasta*.

Sea Bream. Fam. SPARIDAE.

„ (Blkr.). *Gerres abbreviatus*.

„ *filamentosus*.

“Silver-Bream.” Fam. GERRIDAE.

Karang. Reef or coral.

Ikan karang. Fish frequenting rocks and coral reefs.

Kasi-kasi. *Engraulis baelama*.

(“Whitebait”) Herring. Fam. CLUPEIDAE.

Kawan-kawan (R. M.). *Dangila burmanica*.

„ *cuvieri*.

Carp. Fam. CYPRINIDAE.

Kēbasi (Pahang) = **Sēlangat** q.v.

Kēdēmut. *Caranx* sp.

Horse-Mackerel. Fam. CARANGIDAE.

Kēdēra. See **Bēlanak**.

Grey Mullet. Fam. MUGILIDAE.

Kēdewas. See **Bawal**.

Pomfret. Fam. STROMATEIDAE.

Kēdondong. A large **bulus-bulus**.

See **Bulus-bulus**.

Whiting. Fam. SILLAGINIDAE.

Kekek gēdabang (R. M.). *Equula edentula*.

"Silver-Bream." Fam. GERRIDAE.

Kēkek labu. *Gazza minuta*.

"Silver-Bream." Fam. GERRIDAE.

Kēkek jawa. *Mene maculata*.

Kēkek gēdabang. *Mene maculata*.

Horse-Mackerel. Fam. CARANGIDAE.

Kēlabau (Wilk. 524). A fresh water fish (Unid.).

(Cf. M. W. & de B. III 129). *Osteochilus kelabau*.

Carp. Fam. CYPRINIDAE.

Kēlah (R. M.). *Barbus kolus*.

" " " *stracheyi*.

Carp. Fam. CYPRINIDAE.

Kēlalawer (Blkr.). *Antennarius hispidus*.

Angler-fish. Fam. ANTENNARIIDAE.

Kēlara (See Wilk. 524). The young of the **sēmbilang**.

Cat-fish. Fam. SILURIDAE.

Kēli (Dun. D. R.). *Clarias magur*.

" (S.). " *teysmanni*.

Cat-fish. Fam. SILURIDAE.

Kēmbal mas. *Thynnus thunnina*.

See **Tongkol**.

Mackerel. Fam. SCOMBRIDAE.

NOTE:—**Kembal mas** and **Tombol mas** derived from Tamil **Kombola mach**.

Kēmbong (Dun.). *Caranx calla*.

Horse-Mackerel. Fam. CARANGIDAE.

" (R M.). *Scomber microlepidotus*.

Mackerel. Fam. SCOMBRIDAE.

Kēmējan. Also **Kēmēnyan**.

Rhynchobatus djeddensis.

Beaked-Rays. Fam. RHINOBATIDAE.

Kěmi. See **Gěmi.**

Kěnděrap. *Bagarius* sp. ?

Cat-fish. Fam. SILURIDAE.

Kěpar. An edible fresh water fish, common in ponds and swamps.

See Bleeker Vol. IX *Polyacanthus hasseltii*.

(Plate only: no description).

Fam. OSPHROMENIDAE.

Kěpau (Dun.). *Catopra fasciata*.

Fresh-water Perch. Fam. NANDIDAE.

Kěpau laut (R. M.). *Glyphidodon notatus*.

Coral-fish. Fam. POMACENTRIDAE.

Kěpayat. (See Wilk. 522). A large fish (unid.).

Cf. M. W. & de B. III 109 **Kapyah**.

Mystacoleucus marginatus.

Carp. Fam. CYPRINIDAE.

Keping (R. M. **Kepang**). *Glyphidodon notatus*.

Coral-fish. Fam. POMACENTRIDAE.

Kěpiyat (M. W. & de B. III 179 **Kepiat**).

Puntius schwanefeldi.

Carp. Fam. CYPRINIDAE.

Kěrai (Dun. **Krai**). *Barbus goniosoma*.

„ (R. M. **Kereh**). „ *neilli*.

„ **kunyet**. „ sp.

„ **jělawat**. „ „

Carp. Fam. CYPRINIDAE.

Kěrapu (Dun.). *Epinephelus tauvina*.

„ „ *Cromileptis altivelis*.

„ „ *Plectropoma maculatum*.

„ (Day I 450, Dun. D. R.). *Epinephelus lanceolatus*.

„ (Blkr. R.). „ *fasciatus*.

„ (Blkr.). „ *boelang*.

„ **karang** (Blkr.). „ *miniatus*.

„ **lumpur** (Blkr.). „ *pantherinus*.

„ **bloso** (Blkr.). „ *corallicola*.

„ **tutul** (Blkr.). „ *merra*.

„ **bebeh** (Blkr.). „ *fuscoguttatus*.

„ „ „ „ *sexfasciatus*.

„ „ „ „ *hoevenii*.

„ **lilin** (R. M.). „ *salmoides*.

„ **sonoh**. „ *Cromileptis altivelis*.

Sea-Perch. Fam. SERRANIDAE.

Kërètang. *Epinephelus pantherinus*.

Sea-Perch. Fam. SERRANIDAE.

Këring, Ikan këring. Lit. dried fish.

Amphisile scutata.

Sea-snipe. Fam. AMPHISILIDAE.

Kërisi (Blkr. Gurisi mejrah). *Synagris taeniopterus*.

” ” ” ” ” *japonicus*.

” ” ” ” ” *tolu*.

Sea-Bream. Fam. SPARIDAE.

Kërisi aji-aji. *Synagris nematopus*.

” **bali** (R. M.). *Scolopsis bilineatus*.

Sea-Bream. Fam. SPARIDAE.

Kerong-kerong also **Mëngkerong**.

” ” (D. R.). *Therapon puta*.

” ” ” ” *quadrilineatus*.

” ” ” ” *theraps*.

” ” (Blkr.). ” *jarbua*.

Snapper. Sub-fam. LUTIANINAE.

” ” (D. R.). *Centrogenys vaigiensis*.

Sea-perch. Fam. SERRANIDAE.

Kërosok (R. M.). *Monacanthus cheirocephalus*.

” **padi** (R. M. **Kerusu padi**). *Monacanthus monoceros*.

“Leather Jackets.” Fam. BALISTIDAE.

Kërtakok (D. R.). *Batrachus grunniens*.

Frog-fishes. Fam. BATRACHIDAE.

Ketang also **Kitang** (Dun, D. R.). *Ephippus argus*.

(D. R.). *Holacanthus annularis*.

Coral fish. Fam. CHAETODONTIDAE.

Ketang (Dun.). *Teuthis virgata*.

” ” ” *stellata*.

” (C.). ” *java*.

” ” ” *concatena*.

” ” ” *dorsalis*.

“Black Trevally.” Fam. TEUTHIDIDAE.

Këtarap (R. M.). *Pseudoscarus ghobban*.

“Parrot-wrasse.” Fam. SCARIDAE.

Kētewas. See **Bawal.**

Kia-kia. See **Yu kia-kia.**

Kubal. *Polynemus* spp.

A name applied to large fish of this family.

Threadfins. Fam. POLYNEMIDAE.

Kuda laut (Dun.). *Hippocampus hystrix*.

Kuda-kuda ayer (D. R.). *Hippocampus guttulatus*.

Sea-Horses. Fam. SYNGNATHIDAE.

Kuku. See **Běngkongkong.**

Kuning-kuning. *Lutianus erythropterus*.

Snappers. Sub-fam. LUTIANINAE.

Kurau. *Polynemus paradiseus*.

„ (C.). „ *indicus*.

„ (R.). „ *sextarius*.

Kurau pipit „ *sextarius*.

Kurau janggut (Dun.). „ *tetradactylus*.

Threadfins. Fam. POLYNEMIDAE.

Lais (M. W. & de B. II 204). *Belodontichthys dinema*.

„ „ *Cryptopterus cryptopterus*.

„ (Dun.). „ *micropus*.

Cat-fish. Fam. SILURIDAE.

Lalang (Dun.). *Crossochilus oblongus*.

„ „ *Rasbora daniconius*.

„ *Chela* spp.

Carp. Fam. CYPRINIDAE.

Lali (M. W. & de B. III 24 **Langli**). *Botia hymenophysa*.

Loach. Fam. COBITIDAE.

Lambai. *Teuthis* sp.

“Black-Trevally.” Fam. TEUTHIDIDAE.

Lampam (M. W. & de B. III 178). *Puntius schwanefeldi*.

(R. M.). *Barbus jerdoni*.

Carp. Fam. CYPRINIDAE.

Lampila (S.) (**Lampile**). *Betta bellica*.

Fam. OSPHROMENIDAE.

See **Bělaga**.

- Landok** (Pahang). *Sparus datnia*.
Sea-Bream. Fam. SPARIDAE.
- Langgai**. *Trichiurus savala*.
"Barracouta." Fam. TRICHIURIDAE.
- Langi**. A term applied to **Těnggiri** of the largest size.
- Lau** (East coast). *Polynemus sextarius*.
Jew-fish. Fam. POLYNEMIDAE.
- Lawang** (C. & S. dict. 171).
Cf. M. W. & de B. II 259. *Pangasius* sp.
" " 271. *Bagarius* sp.
Cat-fish. Fam. SILURIDAE.
- Lawi ayam**. See **Bulu ayam**.
- Layer, Layeran** or **Sělayer** (Dun. D. **laiar**). *Histiophorus gladius*.
Sail-fish. Fam. HISTIOPHORIDAE.
- Layur** (D. R.) also **Sělayur**. *Trichiurus savala*.
"Barracouta." Fam. TRICHIURIDAE.
- Lěbai** (R. **Lebis**). *Mulloidides flavolineatus*.
Upeneus luteus.
" *tragula*.
Red-Mullet. Fam. MULLIDAE.
- Lebam**. See **Debam**.
- Lele** (Wilk. 629) Jav. *Clarias punctatus*.
(M. W. & de B. II 189). " *melanoderma*.
" " 191). " *batrachus*.
Cat-fish. Fam. SILURIDAE.
- Lembat** (M. W. & de B. II 190). *Clarias nieuhofi*.
Cat-fish. Fam. SILURIDAE.
- Lěmbu** (Dun.). *Ostracion cornutus*.
Box-fishes. Fam. OSTRACIONTIDAE.
Triacanthus sp.
"Leather-jackets." Fam. BALISTIDAE.
- Lěpu** (Dun.). *Antennarius hispidus*.
Angler-fish. Fam. ANTENNARIIDAE.
- " (Dun. D. R.). *Synancidium horridum*.
" (R.). *Scorpaena polyprion*.
" " *Pterois antennata*.
" **panjang** (R. M.). *Pelor didactylum*.
Goblin-fishes. Fam. SCORPAENIDAE.
Malay varieties are **Lěpu sěmaram**.
" **běranyut**.
" **landak**.

Mata lalat (Hanitsch Jour. S. B. R. A. S. Dec., 1912).

Haplochilus panchax.

"Millions." Fam. CYPRINODONTIDAE.

Lidah also **Lidah-lidah** (Dun.). *Cynoglossus lida.*

" " " *grandisquamis.*

" " " *lingua.*

(Dun. C.). *Plagusia bilineata.*

" *Cynoglossus elongatus.*

" *Psettodes erumei.*

" **baji** (D. R.). *Synaptura orientalis.*

" **lumpur** (D. R.). *Synaptura commersoniana.*

" " (C.). *Cynoglossus cantoris.*

Flat-fish. Fam. PLEURONECTIDAE.

See also **Sa-bĕlah.**

Lisah (C.). *Periophthalmus schlosseri.*

(Mud-Skipper). Goby. Fam. GOBIIDAE.

Logu (D. R.). *Choerops oligacanthus.*

"Parrot-fishes." Fam. LABRIDAE.

" (R. M.). *Myripristis murdjan.*

"Silver-Bream." Fam. BERYCIDAE.

Loma (R. M.). *Thynnichthys sandkhol.*

Carp. Fam. CYPRINIDAE.

Luding. A term applied to small **Tĕnggiri.**

Luli (C.). *Harpodon nehereus.*

See **Lumi.** Fam. SCOPELIDAE.

Lumban (R.). *Teuthis java.*

The "Black Trevally" of Australia.

"Black Trevally." Fam. TEUTHIDIDAE.

Lumi. *Harpodon nehereus.*

The "Bombay-duck." Fam. SCOPELIDAE.

Lundu (M. W. & de B. II 345). *Macrones gulio.*

Cat-fish. Fam. SILURIDAE.

Malong (Dun. D. R.). *Muraenesox talabon.*

" *talabonoides.*

" *cinereus.*

Conger eels. Fam. MURAENIDAE.

Mamong. *Caranx* sp.

Horse-Mackerel. Fam. CARANGIDAE.

- Mandi abu.** *Diagramma* spp.
 “Grunters.” Fam. PRISTIPOMATIDAE.
- „ *Novacula* spp.
 “Parrot-fishes.” Fam. LABRIDAE.
- Mata bëliak** (M. W. & de B. II 68 **Mata belo**).
Clupea (Alosa) kanagurta.
 See **Bëliak mata**.
 Herring. Fam. CLUPEIDAE.
- Mayong.** *Arius* sp.
 Cat-fish. Fam. SILURIDAE.
- Mëmpinis.** *Engraulis* spp.
 (“White-bait.”) Herring. Fam. CLUPEIDAE.
- Mëmpurong.** Also **Porong** or **Purong**.
Lycothrissa crocodilus.
 (Sprat or Anchovy.) Herring. Fam. CLUPEIDAE.
- Mënangin.** *Elops hawaiiensis*.
 Giant Herring. Fam. ELOPSIDAE.
- Mëngkai** or **Mingkai** (Wilk. 651). A species of Ray.
- Mëngkerong.** See *Kerong-kerong*.
- Merah** (R. M.). *Lutianus roseus*.
 Snapper. Sub-fam. LUTIANINAE.
- Merah China.** *Caesio pinjalu*.
 Sea-Bream. Fam. SPARIDAE.
- Mërawan.** *Lutianus* sp.
 Snapper. Sub-fam. LUTIANINAE.
- Mudin** or **Mudim.** *Saurus myops*.
- Mësikor.** *Diagramma* spp.
 “Grunters.” Fam. PRISTIPOMATIDAE.
- „ *Novacula* sp.
 Parrot-fishes. Fam. LABRIDAE.
- Mersuji.** *Histiophorus* sp.
 Said to be smaller than **Sëlayer**.
 Sail-fish. Fam. HISTIOPHORIDAE.
 Fam. SCOPELIDAE.
- Nandong** (Kedah) = **Sëlangat**.
 Herring. Fam. CLUPEIDAE.

Nyior-nyior (D. R. Nior-nior). *Trachynotus ovatus*.

The Dart of Australia.

Horse-Mackerel. Fam. CARANGIDAE.

Nyua-nyua (Dun.). *Barilius guttatus*.

Luciosoma setigerum.

Carp. Fam. CYPRINIDAE.

Otek (Blkr.). *Arius utik*.

Cat-fish. Fam. SILURIDAE.

Pachal. See **Parang-parang**.

Paku. See **Rěnnnyau**.

Pala (Dun.). *Betta pugnax*.

Fam. OSPHROMENIDAE.

See *Bělaga*.

Parang-parang (M. W. & de B. II 18). *Chirocentrus dorab*.

The terms used to describe different sizes of this fish are:

Pachal, largest.

Těgap, large.

Chabok, medium.

Sudip, small.

Chabok sětu or **sětul** is the term applied to this fish when caught, (usually in seine nets) in shallow water: amongst the marine plant (**sětul**).

The Dorab. Fam. CLUPEIDAE.

Pari kěbas (C.). *Narcine timlei*.

„ „ „ *Astrape dipterygia*.

„ **linchin** „ *Temera hardwickii*.

„ **bunga** (C. **bang**a). *Astrape dipterygia*.

Electric-Rays. Fam. TORPEDINIDAE.

Pari dědap (Dun.). *Urogymnus asperrimus*.

„ **běting** „ *Trygon uarnak*.

„ **běndera** „ „ *sephen*.

„ **daun** „ „ „

„ **lalat**. *Trygon walga*.

„ **rěnnnyau** „ *kuhlii*.

Sting-Rays. Fam. TRYGONIDAE.

- Pari kělaware** (C.). *Myliobatis vespertilio*,
 „ „ „ „ *nieuhofti*.
 „ **lang** (C. D.). *Aetobatis narinari*.
 „ **daun** (C.). *Rhinoptera adspersa*.
 „ **paus** (D.). *Dicerobatis eregoodoo*.
 „ „ (Dun.). *Ceratoptera ehrenbergii*.
 „ **kola** „ „ „
 Eagle-Rays. Fam. MYLIOBATIDAE.
- Pasir** (Dun.). *Acanthopsis choerorhynchus*.
 Loaches. Fam. COBITIDAE.
 „ (R. M.). *Labeo boggut*.
 Carp. Fam. CYPRINIDAE.
- Pasir-pasir** also **Měmpasir** (Blkr.). *Scolopsis personatus*.
 „ „ *cancellatus*.
 „ „ *ciliatus*.
 „ „ *bilineatus*.
 „ „ *vosmaeri*.
 „ „ *bimaculatus*.
 „ „ *monogramma*.
 Sea-Bream. Fam. SPARIDAE.
- Patin** (M. W. & de B. II 257). *Pangasius pangasius*.
 Cat-fish. Fam. SILURIDAE.
- Patong** (S. **petong**). *Catopra fasciata*.
 Fresh-water Perch. Fam. NANDIDAE.
- Pěchah pěriok** (C.). *Lobotes surinamensis*.
 The Dusky Perch. Fam. LOBOTIDAE.
- Pědukang** (M. W. & de B. II 327). **Anak dukang**.
Hemipimelodus borneensis.
 Cat-fish. Fam. SILURIDAE.
 See **Bělukang**.
- Pělaga**. Also **Ampit-ampit**, **Pala** and **Bělaga**.
Betta pugnax.
 „ *bellica*.
 Fam. OSPHROMENIDAE.
- Pělaling**. Siamese, pla = fish.
 Horse mackerel. Fam. CARANGIDAE.
- Pělata** (Meek, Siamese Pla-thu). *Scomber microlepidotus*.
 Mackerel. Fam. SCOMBRIDAE.
 Varieties are **Pělata Bali** and **Pělata minyak**.

Petek-petek (Dun.). *Ambassis ranga*.

” “ ” ” ” *commersonii*.

Sea-Perch. Fam. SERRANIDAE. Sub-fam. AMBASSINAE.

Pěti, Ikan pěti. A name applied to the Box fishes.

See Buntal. Fam. OSTRACIONTIDAE.

Petong (S.). *Catopra fasciata*.

Fresh-water Perch. Fam. NANDIDAE.

Pias. *Dorosoma chacunda*.

Herring. Fam. CLUPEIDAE.

Pijat-pijat. *Scolopsis torquatus*.

Sea-Bream. Fam. SPARIDAE.

Pinang-pinang (D. R.). *Chaetodon octofasciatus*.

(R. M.). ” *vagabundus*.

Coral fish. Fam. CHAETODONTIDAE.

Sparus datnia.

Sea-Bream. Fam. SPARIDAE.

Pipit (D. R.). *Chelmo rostratus*.

Coral fish. Fam. CHAETODONTIDAE.

Ponggok. A fish inhabiting reefs. Unidentified.

Porong. See mēmpurong.

Puchuk (C.). *Trichiurus savala*.

” *haumela*.

“ Barracouta.” Fam. TRICHIURIDAE.

Puchok pisang (Unid.). Carp. Fam. CYPRINIDAE.

Pukul gendang (R. M.). *Percis pulchella*.

Star-gazer. Fam. LEPTOSCOPIDAE.

Puntong damar. See Bēlanak.

Puput. Also **Puput Malacca**.

” (M. W. & de B. II 90). *Pellona amblyuropterus*.

” *elongata*.

(M. W. & de B. II 93) ” *dussumieri*.

Raonda russelliana.

Herring. Fam. CLUPEIDAE.

Puput (R.). *Hemirhamphus limbatus*.

Puput banang ” *far*.

Gar-fish. Fam. SCOMBRESOCIDAE.

- Puteh** (Dun. as *Barbus maculatus*). *Puntius binotatus*.
 „ („ „ „ *apogon*). *Cyclocheilichthys apogon*.
 „ („). *Rasbora vulgaris*.
 „ („ as *Barbus obtusirostris*). *Mystacoleucus marginatus*.
 Carp. Fam. CYPRINIDAE.
- Puyu.** Also **Puyu-puyu** and **Pěpuyu**.
 „ (S. Dun). *Anabas scandens*.
 The Climbing Perch. Fam. OSPHROMENIDAE.
- Rapang.** Also **Rěpang**.
 See **Bělanak rapang** and **Gělama rapang**.
- Rěnniau.** *Atherina forskali*.
 „ *temmincki*.
 Sand-Smelts. Fam. ATHERINIDAE.
- Riu-riu.** (Dun.). *Lais heranema*.
 Cat-fish. Fam. SILURIDAE.
- Rong** (Dun.). *Dangila cuvieri*.
 „ **běras** (C. & S. dict. 271). Idem ?
 „ (R. M.). *Labeo caeruleus*.
 Carp. Fam. CYPRINIDAE.
- Rumbong-rumbong** (R. M.). *Lutianus madras*.
 Snapper. Sub-fam. LUTIANINAE.
- Rumi-rumi** (D. R.). *Echineis naucrates*.
 Sucking-fish. Fam. ECHINEIDIDAE.
- Sa-bělah** (Dun.). *Synaptura achira*.
 (C.). *Psettodes erumei*.
 „ *Pseudorhombus russellii*.
 Flat-fish. Fam. PLEURONECTIDAE.
 See **Lidah**.
- Sagai** (R.). *Caranx gallus*.
 „ (D. R.). „ *armatus*.
 Horse-Mackerel. Fam. CARANGIDAE.
- Sai** (Wilk. 367). A kind of Ray.
- Sěbarau** also **Barau-barau** (Dun.). *Hampala macrolepidota*
 (*Barbus hampal.*)
 (R. M.). „ *hexastichus*.
 Carp. Fam. CYPRINIDAE.
- Sěběkah.** *Apogon* spp.
 Sea-Perch. Sub-fam. CHILODIPTERINAE.

Sěběkah karang. *Myripristis murdjan.*

Nannygai. Fam. BERYCIDAE.

Sědakang (R. M.). *Gerres altispinnis.*

"Silver-Bream. Fam. GERRIDAE.

Sěkiki. See **Kekek.**

Sělampai (C.). *Collichthys biaurita.*

Jew-fish. Fam. SCIAENIDAE.

Sělangat (M. W. & de B. II 26 **selangkat**).

Dorosoma chacunda.

" **bělau** " *nasus.*

" **tuli** " *sp.*

Herring. Fam. CLUPEIDAE.

Sělangin (C.). *Polynemus tetradactylus.*

" " " *sextarius.*

Thread-fins. Fam. POLYNEMIDAE.

Sělar (Dun). *Caranx kurra.*

" " *Trachynotus bailloni.*

" **batang** (R.). *Caranx djeddaba.*

" **abu-abu** (R. M.). " *ire.*

" **kuning** " " *gymnostethoides.*

" **letup-letup.** " *oblongus.*

" " " *compressus.*

The Malays of Singapore differentiate between three sizes of **Sělar batang**, viz.,

Small, **Sělar renchih.**

Medium, " **kěledek.**

Large, " **batang.**

Other varieties, **Sělar bulat** and **Sělar lepir.**

Horse Mackerel. Fam. CARANGIDAE.

Sělayer. *Histiophorus gladius.*

Sail-fish. Fam. HISTIOPHORIDAE.

Sělayur. *Trichiurus savala.*

"Barracouta." Fam. TRICHTURIDAE.

Sělěmah. *Lactarius delicatulus.*

Fam. LACTARIIDAE.

Sěliap (D. R. **Saliup**). *Chorinemus lysan.*

" " " *sancti-petri.*

Horse-mackerel. Fam. CARANGIDAE.

- Sēlichin.** *Anampses caeruleopunctatus*.
Parrot-fish. Fam. LABRIDAE.
- Sēligi (R.).** *Anacanthus barbatus*.
Leather-jackets. Fam. BALISTIDAE.
- Sēlikor.** The synonym in Singapore for a large **Chēncharu**.
Caranx rotterli.
- Sēlimang** (M. W. & de B. III 230). *Epalzeorhynchus kallopterus*.
Carp. Fam. CYPRINIDAE.
- Sēlinching.** *Pentapus* spp.
"Grunters." Fam. PRISTIPOMATIDAE.
- Sēluang** (S.). *Rasbora argyrotaenia*.
" (Dun.). „ *trilineata*.
" (R.). *Barilius guttatus*.
Small varieties are known as **Sēluwang bēras** C. & S.
dict. 271.
Carp. Fam. CYPRINIDAE.
- Sēludu** (Blkr. as *Arius maculatus*). *Pseudarius arius*.
" (C. **Surdudu**). *Arius macronotacanthus*.
Cat-fish. Fam. SILURIDAE.
- Sēmangka** (D.). *Apogon frenatus*.
Sea-Perch. Sub-fam. CHILODIPTERINAE.
See **Sēbēkah**.
- Sēmaram.** *Centrogenys waigiensis*.
Sea-Perch. Fam. SERRANIDAE.
" (R. M.). *Centropogon indicus*.
" **karang** (R. M.). *Synancia verrucosa*.
"Goblin-fish." Fam. SCORPAENIDAE.
- Sēmbak.** See **Tongkol**.
- Sēmbilang** (D. R.). *Plotosus canius*.
" (C. Blkr.). *Paraplotosus albilabris*.
" **karang** (M. W. & de B. II 230). *Plotosus anguillaris*.
Cat-fish. Fam. SILURIDAE.
- Sēmpila.** Also **Sēmpilai** (Wilk.) and **Lampile** (S.).
See **Bēlaga**.
Fighting-fish. Fam. OSPHROMENIDAE.
- Sēnangin** (C.). *Polynemus tetradactylus*.
" „ *sextarius*.
Thread-fins. Fam. POLYNEMIDAE.

Sëndarat. *Lutianus argentimaculatus*.

Snapper. Sub-fam. LUTIANINAE.

Sënderong (D. **senderang sendok**). *Plectropoma maculatum*.
Epinephelus sexfasciatus.

Sea-Perch. Fam. SERRANIDAE.

Sëngaring. See Wilk. 384. Also **Karing**.

Cf. M. W. & de B. III 152. **Sëngkaring**.

Labeobarbus tambra.

Carp. Fam. CYPRINIDAE.

Sënohong. A large **Sënangin**.

Sënyor. *Psettus falciformes*.

Bat-fish. Fam. SCORPIDIDAE.

Sëpat. (Dun). *Osphromenus trichopterus*.

Fam. OSPHIROMENIDAE.

Sëpat karang (R. M.). *Lobotes surinamensis*.

The Dusky Perch. Fam. LOBOTIDAE.

Sëpat karang. *Pempheris* spp.

"Bull's Eyes." Fam. PEMPHERIDAE.

Sërandong (See Wilk. 381). A fresh water fish. (Unidentified).

"It is something like the **Sëlangat**."

Sërasah. Literally rubbish, manure.

Ikan sërasah. Small and immature fish used as manure.

Sëriding (Dun.). *Equula edentula*.

"Silver-Bream." Fam. GERRIDAE.

Sëtoka. A small Ray.

Sëtonggang. *Monocentris japonicus*.

Knight-fishes. Fam. MONOCENTRIDAE.

Sëtu or **Sëtul**. A marine plant with edible fruit. *Enhalus acoroides*.

See **Parang-parang sëtu** and **Tamban sëtu**.

Sia-sia (R. M.). *Diploprion bifasciatum*.

Ambassis gynocephalus.

Sea-Perch. Fam. SERRANIDAE.

Siakap (C.). *Lates calcarifer*.

Sea-Perch. Fam. SERRANIDAE.

Sirat-sirat. A marine eel (unid.).

Songsong arus. *Caranx* sp.

Horse-mackerel. Fam. CARANGIDAE.

Sudip. Anak sudip. The young of the **Ikan parang.**

Sumpit. Sumpit-sumpit.

(Dun. C. D. R.). *Toxotes chatareus*.

(Blkr.). „ *jaculator*.

Blow-pipe fish. Fam. TOXOTIDAE.

„ *Chelmo rostratus*.

Chaetodon vagabundus.

Coral-fish. Fam. CHAETODONTIDAE.

Susur batang (Dun. **Sulir batang**, as *Rasbora daniconius*.)

Rasbora einthoveni.

Carp. Fam. CYPRINIDAE.

Talang (Dun.). *Chorinemus lysan*.

„ *sancti-petri*.

„ *moadetta*.

Large specimens of *Chorinemus* are usually called

Talang and small ones **Sēliap**.

Horse-mackerel. Fam. CARANGIDAE.

Tali (S.) ? *Acanthopsis choerorhynchus*.

Loach. Fam. COBITIDAE.

Tambak. See **Bawal tambak**.

Tamban (M. W. & de B. II 58 **tembang**). *Clupeoides lile*.

„ (M. W. & de B. II 76). *Clupea* (*Harengula*) *fimbriata*.

„ **bětul** (C. **batal**). *Clupea perforata*.

„ **bulat** (C.). *Dussumieria acuta*.

„ **nipis** (C.). *Clupea perforata*.

„ **siantan**. *Clupea* (*Harengula*) *fimbriata*.

„ „ „ (*Amblygaster*) *leiogaster*.

„ **chinchang rēbong**. *Clupeoides lile*.

„ **jěboh**. *Dussumieria acuta*.

„ „ „ *hasseltii*.

Tamban. Anak tamban jěboh. *Spratelloides delicatulus*.

„ *gracilis*.

Also **Tamban sětu** (unid.) & **Tamban běluru** (unid.)

(“Sardines”) Herring. Fam. CLUPEIDAE.

Tamběra. Also **Těmběra** or **Těběra**.

„ (M. W. & de B. III 152). *Labeobarbus tambra*.

Carp. Fam. CYPRINIDAE.

Tampok-tampok (Wilk. 187). *Gerres oblongus*.

“Silver-Bream.” Fam. GERRIDAE.

Tanda-tanda (Wilk. 193). *Lutianus sillao*.

„ *bohar*.

Snapper. Sub-fam. LUTIANINAE.

Tapa also **Tapah** (Dun.). *Callichrous pabda*.

Cf. M. W. & de B. II 202. *Wallago* spp.

Cat-fish. Fam. SILURIDAE.

Těbal bibir (R. M.). *Diagramma cinctum*.

„ „ „ „ *punctatum*.

„ „ (D. R.). „ *crassipinum*.

„ „ (R.). „ *pictum*.

“Grunters.” Fam. PRISTIPOMATIDAE.

Těbal pipi = **Gěrut-gěrut** q.v.

Těkok (onom.). *Haileutaea stellata*.

Croakers. Fam. MALTHIDAE.

Telan. See **Tilan**.

Tělan rumput (R. M.). *Kyphosus (Pimelepterus) cinerascens*.

“Drummer.” Fam. KYPHOSIDAE.

Těmbakul. *Periophthalmus schlosseri*.

(Mud-skipper) Goby. Fam. GOBIIDAE.

Těmbělian. *Barbus* sp.

Carp. Fam. CYPRINIDAE.

Těmběreh (C. Tembari). *Sciaena diacanthus*.

(Wilk. 181). *Otolithus punctatus*.

Jew-fish. Fam. SCIAENIDAE.

Těměngalan (R. M.). *Barbus burmanicus*.

(Blkr. **Teban-galang**). *Amblyrhynchichthys truncatus*.

Carp. Fam. CYPRINIDAE.

Těměnggong (Blkr.). *Priacanthus tayenus*.

Sea-perch. Fam. SERRANIDAE.

Těměnong = **Kěmbong**. q.v.

Těmoleh (R. **Tamaleh**, as *Barbus apogon*.) *Cyclocheilichthys apogon*.

Carp. Fam. CYPRINIDAE.

Těmpěras (R. M. as *Barbus apogon*.) *Cyclocheilichthys apogon*.

Těmpuwa (Wilk. 188 as *Barbus apogon*.) *Cyclocheilichthys apogon*.

Carp. Fam. CYPRINIDAE.

Těnggiri (Dun.). *Cybium kuhlii*.

„ (C.). „ *commersonii*.

„ „ „ *lineolatum*.

„ **batang** (D.). „ *commersonii*.

„ **musang** „ „

„ **papan** (C.). „ *guttatum*.

The descriptive terms **Těnggiri luding**, **T. tohok**, **T. padi**, **T. tundan** and **Langi** are used with reference to the size of these fish.

Mackerel. Fam. SCOMBRIDAE.

The **Těnggiri** is the well-known sporting fish, the Spanish Mackerel of the Philippines and Australia.

Těngkerong. See **Kerong-kerong**.

Těnok. *Sphyræna novae-hollandiae*.

„ *obtusata*.

„ *jello*.

Small **Kachang-kachang**, larger **Alu-alu**, largest size

Těnok.

“Barracudas.” Fam. SPHYRAENIDAE.

Těrbul. *Osteochilus hasselti*.

Cf. S. “**Teboye**” Duncker p. 205.

Carp. Fam. CYPRINIDAE.

Těri (M. W. & de B. II 46). *Stolephorus commersonii*.

„ „ „ *indicus*.

„ „ „ *tri*.

(“Whitebait”) Herring. Fam. CLUPEIDAE.

Těripang (R. M.). *Saurus indicus*.

Fam. SCOPELIDAE.

Těrubok (M. W. & de B. II 66). *Clupea (Alosa) macrura*.

„ **padi** „ „ „ *toli*.

„ **korin** „ „ „ *sp*.

Herring. Fam. CLUPEIDAE.

Tilan. Also **telan** (Dun.). *Mastacembulus unicolor*.

„ „ „ *maculatus*.

(Dun. D.). „ *armatus*.

Spiny-Eels. Fam. MASTACEMBELIDAE.

Tilan pasir (D. R.). *Trypauchen vagina*.

Goby. Fam. GOBIIDAE.

Timah-timah. See also **Sĕlayur** and **Langgai**.

„ (Dun.). *Trichiurus savala*.

„ „ *haumela*.

“Barracouta.” Fam. TRICHIURIDAE.

Timun-timun also **Mĕntimun** (R. M.). *Lutianus lineolatus*.

„ *decussatus*.

„ *vitta*.

Snapper. Sub-fam. LUTIANINAE.

Tiram. *Engraulis* sp.

Herring. Fam. CLUPEIDAE.

Todak, also **Chĕnchodak** (Dun.). *Belone cancila*.

„ (Dun. C. R.). „ *strongylurus*.

„ (Dun. D.). „ *choram*.

„ (C.). „ *annulata*.

„ **pendek** (Penang) (Blkr.). *Hemirhamphus georgii*.

„ „ (Malacca) „ „ *far*.

Gar-fish. Fam. SCOMBRESOCIDAE.

Toka-toka also **Sĕtoka**. A small Ray.

Tokak (Blkr. **Toka**). *Cossyphus diana*.

„ *Chaerops omnopterus*.

Parrot-fish. Fam. LABRIDAE.

Toman (R.). *Ophiocephalus striatus*.

“Murrel.” Fam. OPHIOCEPHALIDAE.

Tombol mas (R. M.). *Thynnus thunnina*.

The Tunny. See **Kĕmbal mas**.

Mackerel. Fam. SCOMBRIDAE.

Tombong damar (Wilk. 181). A fish (unid.)

See **Puntong damar**.

Tongkol. *Thynnus thunnina*.

Small sized fish are called **Choreng**, medium sized specimens **Sĕmbak**, large ones **Tongkol** and exceptionally large ones **Kĕmbal mas** or **Tombol mas**.

The **Tunny** or **Tuna**.

Mackerel. Fam. SCOMBRIDAE.

Tudong pĕriok also **Tudong tĕmpayan**.

(Blkr. **Tudjong-prio**). *Platax batavianus*.

„ *vespertilio*.

Coral-fish. Fam. CHAETODONTIDAE.

Tuli. Literally deaf. See **Sĕlangat tuli.**

Tumbok banir. *Histiophorus* sp.

Sail-fish. Fam. HISTIOPHORIDAE.

Tumbok tĕbing (Dun, **Tumbok ka tubing**). Neg. Semb.

Luciocephalus pulcher.

Fam. OSPHIROMENIDAE.

Tunjang langit. *Triacanthus oxycephalus.*

"Leather-jackets." Fam. BALISTIDAE.

Ubi (C.). *Sillago sihama.*

"Whiting." Fam. SILLAGINIDAE.

Udip. Petit ikan parang (Favre). See **Sudip.**

Umbut-umbut (S. Mombu). *Dangila lineata.*

" (M. W. & de B. III 116 **Umbu-umbu**). *D. cuvieri.*

Barynotus microlepis.

Carp. Fam. CYPRINIDAE.

Undok-undok. *Hippocampus* sp.

The Sea-Horse.

Fam. SYNGNATHIDAE.

Ungar (Wilk. 51). *Lutianus argentimaculatus.*

" *johnii.*

Snapper. Sub-fam. LUTIANINAE.

Unsat or **usat.** *Plotosus* sp.

Cat-fish. Fam. SILURIDAE.

Yu (R. M.). *Carcharias dussumieri.*

" **tĕnggiri.** *Galeocerdo rayneri.*

Sharks. Fam. CARCHARIDAE.

Yu bĕngkong (D.). *Sphyrna (Zygaena) malleus.*

" (D.). " *blochii.*

Hammer-head Sharks. Fam. SPHYRNIDAE.

Yu pendek (D.). *Scyllium marmoratum.*

" **chechak** (D. **Chikak**). *Stegostoma tigrinum.*

" **tokek** (C. **Tokay**). " "

" **tokek** (C. **Tokay**). *Chiloscyllium indicum.*

" **bĕlangkas** (D.). " "

Dog-fishes. Fam. SCYLLIDAE.

" **parang** (Dun.). *Pristis cuspidatus.*

" **gergaji** " " "

" **todak** (D.). " "

Saw-fishes. Fam. PRISTIDAE.

" **kia-kia** (D.). *Rhynchobatus djeddensis.*

" " (D.). *Rhinobatus thonini.*

Beaked-Rays. Fam. RHINOBATIDAE.

Families of Malayan Fishes.

PART III.

ELOPSIDAE (GIANT-HERRINGS).

Elops hawaiiensis Regan.	<i>Bandang, Měnangin.</i>
Megalops cyprinoides Brouss.	<i>Bulan-bulan.</i>

NOTOPTERIDAE (FEATHER-BACKS).

Notopterus notopterus Pall.	<i>Bēlida.</i>
„ chitala H. B.	„

CHANIDAE (THE MILK-FISH).

Chanos chanos Forsk.	<i>Bandang, Jangas.</i>
-----------------------------	-------------------------

CLUPEIDAE (HERRINGS, SHADS, ETC.).

Chirocentrus dorab Forsk.	<i>Parang-parang.</i>
Spratelloides delicatulus Benn.	<i>Anak tamban jěboh.</i>
„ gracilis Schleg.	„ „ „
Dussumieria acuta C. V.	<i>Tamban bulat, T. jěboh.</i>
„ hasseltii Blkr.	<i>Tamban jěboh.</i>
Dorosoma nasus Bl.	<i>Sělangat bėlau, Nandong,</i> <i>Kėbasi.</i>
„ chacunda H. B.	<i>Sělangat, Kėbasi, Nandong,</i> <i>Pias.</i>
Setipinna breviceps Cant.	<i>Biang-biang.</i>
„ taty C. V.	„
Lycothrissa crocodilus Blkr.	<i>Měmpurong.</i>
Engraulis baelama Forsk.	<i>Kasi-kasi.</i>
„ grayi Blkr.	<i>Bangkok.</i>
„ mystax Bl. Schn.	<i>Bulu ayam.</i>
„ setirostris Brouss.	<i>Bangkok.</i>
„ spp.	<i>Tiram, Měmpinis, Jěmbėdi.</i>
Stolephorus commersonii Lac.	<i>Bilis, Tėri.</i>
„ indicus v. H.	<i>Bunga ayer, Tėri.</i>
„ tri Blkr.	<i>Tėri, Bilis.</i>
Coilia dussumieri C. V.	<i>Bulu ayam.</i>
„ quadrifilis Günth.	„ „
Clupeoides lile C. V.	<i>Tamban, T. chinchang rė</i> <i>bong.</i>

Clupea (Amblygaster) clupeoides	Blkr.	<i>Tamban.</i>
„ (Amblygaster) leiogaster	C. V.	<i>Tamban siantan.</i>
Clupea (Alosa) toli	C. V.	<i>Tërubok padi.</i>
„ „ macrura	Blkr.	<i>Tërubok, Ikan bengkalis.</i>
„ „ kanagurta	Blkr.	<i>Bèliak mata.</i>
„ „	spp.	<i>Tërubok korin.</i>
„ (Harengula) fimbriata	C. V.	<i>Tamban, T. siantan.</i>
„ „ moluccensis	Blkr.	<i>Bèliak mata jantan.</i>
„ „ perforata	Cant.	<i>Tamban bêtul, T. nipis.</i>
Pellona amblyropterus	Blkr.	<i>Puput, P. Mèlaka.</i>
„ elongata	Benn.	„ „
„ dussumieri	C. V.	„ „ <i>Bèliak mata, kapak.</i>
Opisthopterus tartoor	C. V.	<i>Chandong.</i>
Raconda russelliana	Gray.	<i>Puput, Chandong.</i>

SCOPELIDAE (QUEENSLAND-SMELT, ETC.).

Saurida tumbil	Bl.	<i>Bèlungkor.</i>
Saurus myops	Bl. Schn.	<i>Mudin or Mudim.</i>
„ indicus	Day.	<i>Tèripang.</i>
Harpodon nehereus	H. B.	<i>Luni, Luli.</i>

CYPRINODONTIDAE (“MILLIONS”).

Haplochilus panchax.	<i>Mata lalat.</i>
-----------------------------	--------------------

SILURIDAE (CAT-FISH).

Clarias melanoderma	Blkr.	<i>Lele.</i>
„ nieuhofi	C. V.	<i>Lembat.</i>
„ batrachus	L.	<i>Lele, Kèli.⁽¹⁾</i>
„ teysmanni	Blkr.	<i>Kèli.</i>
Silurichthys phaiosoma	Blkr.	<i>Gemang darat.</i>
Wallago	sp.	<i>Tapu, Tapah.</i>
Belodontichthys dinema	Blkr.	<i>Lais, Bègahak.</i>
Callichrous pabda	H. B.	<i>Tapu, Tapah.</i>
Cryptopterus cryptopterus	Blkr.	<i>Lais.</i>
„ micropus	Blkr.	„
Paraplotosus albilabris	C. V.	<i>Sèmbilang.</i>
Plotosus	sp.	<i>Usat or Usat.</i>
„ canius	H. B.	<i>Sèmbilang, Kèlara, Gemang.</i>
„ anguillaris	Bl.	<i>Sèmbilang karang.</i>

I. Duncker and Rowell give *C. magur* H. B. for Kèli, which is now regarded by Max Weber and de Beaufort as a synonym of *C. batrachus*.

Lais hexanema Blkr.	<i>Riu-riu.</i> ?
Pangasius spp.	<i>Lawang.</i>
" pangasius H. B.	<i>Patin.</i>
" polyuranodon Blkr.	<i>Juara.</i>
Arius thalassinus Rüpp.	<i>Jahan.</i>
" sagor H. B.	<i>Duri.</i>
" leiotetocephalus Blkr.	<i>Pědukang, Bělukang.</i>
" macronotacanthus Blkr.	<i>Sěludu.</i>
" utik Blkr.	<i>Otek.</i>
" maculatus Thunb.	<i>Sěludu.</i>
" spp.	<i>Mayong, Bagok.</i>
Hemipimelodus borneensis Blkr.	<i>Pědukang.</i>
Macrones nigriceps C. V.	<i>Baung.</i>
" nemurus C. V.	<i>Duri, Baung, Engor-engor.</i>
" planiceps C. V.	<i>Baung kuning.</i>
" gulio H. B.	<i>Lundu.</i>
" bleekeri.	<i>Engor-engor.</i>
Bagarius sp. ?	<i>Kěnděrap.</i>

COBITIDAE AND CYPRINIDAE (LOACHES AND CARPS).

COBITIDAE (LOACHES). ..

Acanthopsis choirorhynchus Blkr.	<i>Pasir.</i>
Botia hymenophysa.	<i>Lali.</i>

CYPRINIDAE (CARPS).

Chela oxygaster C. V.	<i>Lalang.</i>
" sp.	"
Rasbora argyrotaenia Blkr.	<i>Sěluang, Chěřěchek.</i>
" trilineata Steind.	<i>Sěluang.</i>
" lateristriata var. sumatrana Blkr.	<i>Puteh.</i>
" einthoveni Blkr.	<i>Susur batang, Lalang.</i>
" vulgaris Duncker.	<i>Puteh.</i>
Luciosoma setigerum C. V.	<i>Nyua-nyua.</i>
Amblyrhynchichthys truncatus Blkr.	<i>Těměngalan.</i>
Mystacoleucus marginatus C. V.	<i>Kěpayat, Puteh.</i>
Dangila cuvieri C. V.	<i>Umbut-umbut, Rong, Kawan-kawan.</i>
" burmanica Day.	<i>Kawan-kawan.</i>
" lincata Sauv.	<i>Umbut-umbut.</i>
Barynotus microlepis Blkr.	" "
Thynnichthys sandkhol Sykes.	<i>Loma.</i>

Osteochilus kelabau Popta.	<i>Kēlabau.</i>
„ hasselti C. V.	<i>Tērbul.</i>
Hampala macrolepidota C. V.	<i>Sēbarau.</i>
Labeobarbus tambra C. V.	<i>Tambēra</i> also <i>Tēmbēra</i> , <i>Garing</i> , <i>Sēngaring.</i>
Cyclocheilichthys apogon C. V.	<i>Bēbēras</i> , <i>Tēmpēras</i> , <i>Chēm-</i> <i>pēras</i> , <i>Puteh</i> , <i>Tēmpua</i> , <i>Tēmoleh.</i>
Puntius schwanefeldi Blkr.	<i>Lampam</i> , <i>Kēpiat.</i>
„ binotatus C. V.	<i>Puteh</i> , <i>Kērai.</i>
Barbichthys laevis C. V.	<i>Bēntulu.</i>
Labeo caeruleus Day.	<i>Rong.</i>
„ boggut Sykes.	<i>Pasir.</i>
Epalzeorhynchus kallopterus Blkr.	<i>Sēlimang.</i>
Crossochilus oblongus C. V.	<i>Lalang.</i>
Barbus burmanicus Day.	<i>Tēmēngalan.</i>
„ jerdoni.	<i>Lampam.</i>
„ hexastichus McLell.	<i>Sēbarau.</i>
„ neilli Day.	<i>Kērai.</i>
„ kolus Blkr.	<i>Kēlah.</i>
„ stracheyi Day.	„
„ oatesii Blgr.	<i>Daun.</i>
Barilius guttatus Day.	<i>Sēluang</i> , <i>Nyua-nyua.</i>
(Unidentified).	<i>Darok-darok</i> , <i>Puchok pisang.</i> <i>Kērai jēlawat</i> , <i>K. kunyet.</i>

ANGUILLIDAE, CONGRIDAE, OPHICHTHYIDAE, ETC.

(EELS, CONGER-EELS, ETC.).

Muraenesox cinereus Forsk.	<i>Maḥong.</i>
„ talabon Cant.	„
„ talabonoides Blkr.	„
Pisoodonophis cancrivorus Rich.	<i>Bēlin.</i>
Muraena (gymnothorax) undu- lata Lac.	„
(Unidentified).	<i>Laki sēmbilang</i> , <i>Sirat-sirat.</i> <i>Bēlidang.</i>

SYMBRANCHIDAE (SWAMP-EELS).

Monopterus albus Zuiew.	<i>Bēlut.</i>
Macrotema caligans Cant.	<i>Balut.</i>

**SYNGNATHIDAE, AMPHISILIDAE (SEA-HORSES
AND SKELETON-FISHES).**

Hippocampus guttulatus Cuv.	<i>Kuda-kuda.</i>
„ hystrix K. P.	<i>Kuda laut.</i>
Amphisile scutata L.	<i>Kěring.</i>

**SCOMBRESOCIDAE (GAR-PIKES, GAR-FISHES AND
FLYING-FISHES).**

Belone cancila H. B.	<i>Today.</i>
„ strongylurus v. H.	„
„ choram Forsk.	„
„ annulata C. V.	„
Hemirhamphus limbatus C. V.	<i>Puput.</i>
„ far Forsk.	<i>Today pendek. Puput ban- ang.</i>
„ cantoris Blkr.	<i>Jolong-jolong, Jěnjolong.</i>
„ buffonis C. V.	„ „
„ pogonognathus Blkr.	„ „
„ fluviatilis Blkr.	„ „
„ georgii C. V.	<i>Today pendek.</i>
Exocoetus oligolepis Blkr.	<i>Bělanak.</i>
„ neglectus Blkr.	„
„ nigripinnis C. V.	„
„ speculiger Val.	„

ATHERINIDAE (SAND-SMELTS).

Atherina forskali.	<i>Rěnnau, Paku.</i>
„ temmincki.	„ „

MUGILIDAE (GREY MULLET).

Mugil planiceps C. V.	<i>Bělanak, Jěmpul.</i>
„ speigleri Blkr.	„
„ vaigiensis Q. G.	„ <i>B. tamok.</i>
„ cunnesius C. V.	„
„ bleekeri ?	<i>Bělanak rapang.</i>
„ oeur Forsk.	„ <i>tamok.</i>
„ borneensis Blkr.	„ <i>anding.</i>
„ spp.	„ <i>angin, B. puteh, B. kě- děra, Puting damar, Puntong damar, Tom- bong damar, B. ba- kau, Pělong.</i>

POLYNEMIDAE (*THREADFINS*).

Polynemus indicus Shaw.	Kurau, K. janggut, Kubal.
„ sextarius Bl. Schn.	Kurau, K. pipit, Lau.
„ tetradactylus Shaw.	Kurau janggut, Sėnangin, Sėnohong, Kubal.
„ paradiseus Bl.	Kurau.

SPHYRAENIDAE (*BARRACUDAS*).

Sphyraena novae-hollandiae Günth.	Tėnok, Alu-alu, Kachang-kachang.
„ obtusata C. V.	„ „ „
„ jello C. V.	„ „ „
„ commersonii C. V.	„ „ „

STROMATEIDAE (*POMFRETS*).

Stromateus atous C. V.	Bawal chėrmin.
„ cinereus Bl.	Bawal puteh, Bawal itam, Bawal kėdewas.
„ niger Bl.	Bawal tambak.

OPHIOCEPHALIDAE (*MURREL*).

Ophiocephalus gachua H. B.	Aruan, Bėrchat.
„ lucius C. V.	„ „
„ striatus Blkr.	„ Toman.
„ spp.	Gabus, Bujok.

BERYCIDAE (*NANNYGAI*).

Myripristis murdjan Forsk.	Sėbėkah karang, Logu.
-----------------------------------	-----------------------

MONOCENTRIDAE (*KNIGHT-FISHES*).

Monocentris japonicus Bl. Schn.	Sėtonggang.
--	-------------

PEMPHERIDAE (*BULL'S-EYES*).

Pempheris mangula C. V.	Sėpat karang.
„ spp.	„ „

KYPHOSIDAE (*DRUMMERS*).

Kyphosus cinerascens Forsk.	Tėlan rumput.
„ sp.	Bėras-bėras.

LOBOTIDAE (*DUSKY-PERCH*).

Lobotes surinamensis Bl.	Pėchah pėriok, Sėpat karang.
---------------------------------	------------------------------

TOXOTIDAE (*BLOW-PIPE FISHES*).

Toxotes jaculator Pall.	Sumpit-sumpit.
„ chatareus H. B.	„

NANDIDAE (FRESH-WATER PERCHES).

Catopra fasciata Blkr. *Kěpau, Petong, Patong.*

SERRANIDAE (SEA PERCHES).**Serraninae.**

Centrogenys vaigiensis Q. G. *Kerong-kerong, also Měng-kerong, Sēmaram.*

Cromileptes altivelis C. V. *Kěrapu, Kěrapu sonoh.*

Plectropoma maculatum C. V. *Kěrapu, Sěnděrong.*

Epinephelus tauvina Forsk. *Kěrapu, K. lilin, K. kayu.*

 " **lanceolatus** Bl. *Kěrapu.*

 " **fasciatus** Forsk. " *Sěnděrong.*

 " **boelang** C. V. " "

 " **miniatus** Forsk. " *karang.*

 " **pantherinus** Blkr. *Kěrětang, Kěrapu lumpur.*

 " **corallicola** Blkr. *Kěrapu.*

 " **merra** Bl. " "

 " **fuscoguttatus** Forsk. " "

 " **hoevenii** Blkr. " "

 " **salmoides** Lac. " *lilin.*

Priacanthinae.

Priacanthus tayenus Rich. *Těměnggong.*

 " **hamrur** C. V. *Barau-barau laut.*

Centropominae.

Lates calcarifer Bl. *Siakap, also Kakap.*

Psammoperca vaigiensis C. V. *Gělam.*

Ambassinae.

Ambassis commersonii C. V. *Petek-petek.*

 " **ranga** H. B. " "

 " **gymnocephalus** Lac. *Sia-sia.*

Chilodipterinae.

Apogon frenatus Blkr. *Sěmangka.*

 " spp. *Sěbėkah.*

Lutianinae (Snappers).

Lutianus roseus Day. *Ikan merah, Jěněhak.*

 " **argentimaculatus** Forsk. *Ungar, Sěndarat.*

 " **lineolatus** Rüpp. *Timun-timun, also Měn-timun.*

 " **johnii** Bl. *Ungar, Jěněhak.*

 " **sebae** C. V. *Jěněhak.*

 " **fulviflamma** Forsk. " "

 " **lioglossus** Blkr. " "

Lutianus erythropterus Bl.	<i>Kuning-kuning.</i>
„ madras C. V.	<i>Rumbong-rumbong.</i>
„ sillao Russell.	<i>Tenda-tanda.</i>
„ bohar Forsk.	„
„ decussatus C. V.	<i>Timun-limun, Měntimun.</i>
„ vitta Q. G.	„
„ spp.	<i>Bambangan, Mambang or Bambang, Měrawan, Běr- kil, Sěnggaral.</i>
Therapon theraps C. V.	<i>Kěrong-kěrong, also Měng- kerong and Těngkerong.</i>
„ jarbua Forsk.	<i>Kěrong-kěrong.</i>
„ puta C. V.	„
„ quadrilineatus Bl.	„
Diploprion bifasciatum K. V. II.	<i>Sia-sia.</i>
Mesoprion sp.	<i>Gěrut-gěrut.</i>

SILLAGINIDAE (“WHITINGS”).

Sillago sihama Forsk.	<i>Ubi, Bulus-bulus, Běbulus, Kědondong.</i>
„ maculatus Q. G.	<i>Ubi, Bulus-bulus, Běbulus, Kědondong.</i>

SCIAENIDAE (JEW-FISHES).

Sciaena diacanthus Lac.	<i>Těmběreh.</i>
Ucbrina russellii C. V.	<i>Gělama.</i>
Otolithus maculatus C. V.	<i>Jarang gigi.</i>
„ argenteus C. V.	<i>Gělama panjang, Jarang gigi.</i>
„ ruber Bl. Schn.	<i>Jarang gigi.</i>
„ spp.	<i>Gělama panjang, G. papan, G. China, G. sěkang or sěngkang, G. rapang, G. batu, G. itam, G. perak, G. batu Kěling, G. lanjut, G. kuning dada, G. dahi tinggi, G. chěrua, G. pi- sang.</i>
Collichthys biaurita Cant.	<i>Sělampai, Jarang gigi.</i>

GERRIDAE (“SILVER-BREAM”).

Gerres filamentosus C. V.	<i>Kapas-kapas.</i>
„ abbreviatus Blkr.	„
„ altispinnis ?	<i>Sědakang.</i>
„ oblongus C. V.	<i>Tampok-tampok.</i>

Equula edentula Bl.	<i>Kekek gēdabang, Sēriding.</i>
Gazza minuta Bl.	<i>labu.</i>

LACTARIIDAE (SELEMAH).

Lactarius delicatulus C. V.	<i>Sēlēmah.</i>
------------------------------------	-----------------

PRISTIPOMATIDAE (GRUNTERS).

Pristipoma maculatum Bl.	<i>Gērut-gērut. Chēlek mata.</i>
hasta Bl.	"
operculare Playfair.	<i>Ampas tēbu.</i>
guoraca C. V.	<i>Gērut-gērut.</i>
Diagramma crassipinum Rüpp.	<i>Tēbal bibir.</i>
pictum Thunb.	" "
cinctum Tem. Schleg.	" "
punctatum Blkr.	" "
spp.	<i>Bēsikor, Mēsikor, Mandi abu, Kachi.</i>
Pentapus caninus Blkr.	<i>Anjang-anjang.</i>
sp.	<i>Sēlinching.</i>

SPARIDAE (SEA-BREAMS).

Scolopsis ghanam Forsk.	<i>Anjang-anjang.</i>
cancellatus C. V.	<i>Pasir-pasir.</i>
ciliatus Lac.	"
vosmeri Bl.	"
bimaculatus C. V.	"
monogramma K. V. H.	"
personatus C. V.	<i>Gērētak lantai.</i>
bilineatus Bl.	<i>Kērisi Bali.</i>
torquatus C. V. ⁽¹⁾	<i>Pijut-pijat.</i>
Synagris notatus Day.	<i>Kērisi.</i>
japonicus Günth.	<i>" , Gērētak lanta.</i>
taeniopterus C. V.	"
tolu C. V.	"
nematopus Blkr.	<i>" aji-aji.</i>
Caesio kuning Bl.	<i>Dēlah.</i>
lunaris Ehr.	"
pinjalu Blkr.	<i>Jalu-jalu, Jēnjalu, Ikan merah China, Dēlah.</i>
chrysozona K. V. H.	<i>Dēlah karang.</i>

1, Day gives *S. torquatus* = *S. vosmeri* the former being the young and the latter the adult, but Bleeker regards them as separate species, as do the Malays.

Proteracanthus sarissophorus	<i>Batu.</i>
Cant.	
Lethrinus nebulosus Forsk.	<i>Asoh-asoh, Gërëtak lantai.</i>
Sparus hasta Bl. Schn. ⁽¹⁾	<i>Kapas-kapas, Běngkong-kong, Běkukong, Kuku, Bandan.</i>
„ datnia H. B.	<i>Pinang-pinang, Lar-dok.</i>

MULLIDAE (RED MULLET).

Upeneus tragula Richardson.	<i>Biji nangka.</i>
„ luteus Blkr.	<i>Lěbai.</i>
Mulloidies flavolineatus Lac.	„

SCORPIDIDAE (BAT-FISHES).

Psettus argenteus L.	<i>Gědabang.</i>
„ falciformis Lac.	<i>Sěnyor.</i>

CHAETODONTIDAE (CORAL-FISHES AND BUTTER FISHES).

Ephippus orbis Bl.	<i>Daun bāharu.</i>
„ argus L.	<i>Ketang.</i>
Chelmo rostratus L.	<i>Pipit, Sumpit-sumpit.</i>
Heniochus macrolepidotus L.	<i>Gombing.</i>
Holacanthus sexstriatus C. V.	<i>Inggu.</i>
„ mesoleucus Bl.	„
„ annularis Bl.	<i>Ketang.</i>
„ spp.	<i>Babi.</i>
Platax teira Forsk.	<i>Daun, Bonang.</i>
„ batavianus C. V.	<i>Tudong pěriok.</i>
„ vespertilio Bl.	„ „
Chaetodon octofasciatus L.	<i>Pinang-pinang.</i>
„ vagabundus L.	<i>Sumpit-sumpit, Pinang-pinang.</i>

DREPANIDAE (MOON-FISH).

Drepane punctata L.	<i>Daun bāharu.</i>
----------------------------	---------------------

I. According to Day, *S. hasta* = *S. berda*.

TEUTHIDIDAE ("BLACK TREVALLY").

Teuthis nebulosa.	<i>Děngkis.</i>
„ virgata C. V.	<i>Ketang, Děngkis.</i>
„ stellata Forsk.	„
„ java L.	<i>Lumban, Ketang, Debam, Lebam.</i>
„ concatena C. V.	<i>Gelibas, Ketang.</i>
„ dorsalis C. V.	<i>Ketang.</i>
„ oramin Günth. ⁽¹⁾	<i>Gelibas, Bělibas.</i>
„ sp.	<i>Lambai.</i>

OSPHROMENIDAE (GOURAMI. FIGHTING FISHES, ETC.).

Osphromenus olfax L.	<i>Kalui.</i>
„ malayanus Duncker.	<i>Biji durian.</i>
„ trichopterus Pall.	<i>Sěpat.</i>
Anabas scandens Dald.	<i>Puyu-puyu, Pěpuyu, Bětok.</i>
Luciocephalus pulcher Gray.	<i>Tumbok těbing.</i>
Betta pugnax Cant.	<i>Ikan bělaga, Sěmpila, Pě-laga, Pala.</i>
„ bellica Sauv.	<i>Ikan bělaga, Sěmpila, Pě-laga, Lampila.</i>
„ ssp.	<i>Anak ampit, A. karing, Karing gajah.</i>
Polyacanthus hasselti C. V.	<i>Kěpar.</i>

POMACENTRIDAE (CORAL-FISHES).

Amphiprion ephippium Bl.	<i>Inngu.</i>
„ frenatus Brev.	„
Dascyllus sp.	„
Pomacentrus albofasciatus Schleg.	„
Glyphidodon coelestinus C. V.	<i>Gombing.</i>
„ notatus Day.	<i>Kěpau laut, Keping.</i>

LABRIDAE ("WRASSES" OR PARROT-FISHES).

Chaerops omnopterus Rich.	<i>Tokak.</i>
„ oligacanthus Blkr.	<i>Logu.</i>
Cossyphus diana Lac.	<i>Tokak.</i>
Chilinus fasciatus Bl.	<i>Bayan, Boyan.</i>
„ chlorurus Bl.	<i>Jampong, Bayan, Boyan.</i>
Anampses coeruleopunctatus Rüpp.	<i>Sělichin.</i>

I. Day suggests that *oramin* may prove to be synonymous with *dorsalis*

PlatyGLOSSUS dussumieri C. V.	<i>Bēlodok karang.</i>
Novacula spp.	<i>Mandi abu, Mēsikor, Kachi.</i>
Julis lunaris.	<i>Bēchok.</i>

SCARIDAE ("PARROT-WRASSES").

Pseudoscarus ghobbam Forsk.	<i>Kētarap.</i>
„ rivulatus C. V.	<i>Kalat.</i>
Pseudodax moluccanus C. V.	<i>Bēchok.</i>

CARANGIDAE ("HORSE-MACKERELS").

Caranx röttleri Bl.	<i>Chēncharu, Sēlikor, Jarujaru.</i>
„ kalla C. V.	<i>Sēlar, Kēmbong.</i>
„ gallus L.	<i>Sagai, Chērmin.</i>
„ armatus Forsk.	<i>Sagai.</i>
„ kurra C. V.	<i>Sēlar, Kēmbong.</i>
„ gymnostethoides Blkr.	<i>Sēlar kuning.</i>
„ djeddaba Forsk.	<i>Sēlar batang.</i>
„ compressus Day.	<i>Sēlar lētup-lētup, Daing bēlang,</i>
„ ire C. V.	<i>Sēlar abu-abu.</i>
„ boops C. V.	<i>Jalu-jalu (?).</i>
„ oblongus C. V.	<i>Sēlar lētup-lētup.</i>
„ speciosus Forsk.	<i>Daing bēlang.</i>
„ spp.	<i>Bērkas, Songsong arus, Sēlar bulat, S. lepir, Bagat, Mamong, Pēlalang, Kēdēmūt, Gērēpoh, Dēmbudok.</i>
Mene maculata Bl. Schn.	<i>Kekek gēdabang, K. Jawa.</i>
Trachynotus ovatus L.	<i>Nyior-nyior.</i>
„ bailloni Lac.	<i>Sēlar.</i>
Chorinemus moadetta C. V.	<i>Talang, Sēliap.</i>
„ lysan Forsk.	„ „
„ Sti Petri C. V.	„ „

SCOMBRIDAE (MACKERELS, TUNNIES, ETC.).

Scomber microlepidotus Rüpp.	<i>Pēlata, Kēmbong.</i>
Thynnus thunnina C. V.	<i>Ikan ayer, Tongkol, Sēmbak, Choreng, Kembal mas, Tombol mas.</i>

Cybium kahlII C. V.	<i>Tenggiri, Luding, Tohok,</i> <i>Langi.</i>
„ commersonii Lac.	„ „ „ „
„ lineolatum Cuv.	„ „ „ „
„ guttatum Bl. Schn.	<i>T. papan</i> „ „ „

TRICHIURIDAE (HAIR-TAILS).

Trichiurus savala Cuv.	<i>Timah-timah, Selayur,</i> <i>Langgai, Puchuk.</i>
„ haumela Forsk.	<i>Timah-timah, Selayur,</i> <i>Langgai, Puchuk.</i>

HISTIOPHORIDAE (SAIL-FISHES).

Histiophorus gladius Brouss.	<i>Selayur, Layer, Layeran.</i>
„ spp.	<i>Tumbok banir, Mersuji.</i>

PLEURONECTIDAE (FLAT FISHES).

Psettodes erumei Bl.	<i>Lidah, Sabalah.</i>
Pseudorhombus russelli Gray.	„ „
Synaptura achira Duncker.	„ „
„ orientalis Bl. Schn.	„ „
„ commersoniana Lac.	„ „
Cynoglossus lida Blkr.	„ „
„ lingua H. B.	„ „
„ elongatus Günth.	„ „
„ hamiltonii Günth.	„ „
„ cantoris Blkr.	„ „
Plagusia bilineata Bl.	„ „

GOBIIDAE (“GOBIES”).

Eleotris butis H. B.	<i>Belontok.</i>
Gobius giuris H. B.	<i>Belodok.</i>
„ sadanundio H. B.	„ <i>kërapu.</i>
„ sp. aff. caninus C. V. (?)	„ <i>lobang.</i>
„ viridipunctatus ?	<i>Belontok.</i>
Periophthalmus schlosseri Pall.	<i>Tembakul, Lisah, Belodok.</i>
„ koelreuteri Pall.	<i>Belachak.</i>
Trypauchen vagina Bl.	<i>Tilan pasir.</i>
Apocryptes lanceolatus Bl. Schn.	<i>Belodok.</i>
Boleophthalmus boddarti Pall.	„

ECHENEIDIDAE (SUCKING-FISHES).

Echineis naucrates.	<i>Gëmi, Gëdëmi, Rumi-rumi.</i>
----------------------------	---------------------------------

SCORPAENIDAE (GOBLIN-FISHES).

Scorpaena polyprion Blkr.	<i>Lēpu, Dēpu, Gēdēmpu.</i>
Pterois antennata Blkr.	„ „ „
Pelor didactylum Pall.	„ „ „
Centropogon indicus Day.	„ <i>sēmaram.</i>
Synancia verrucosa Bl. Schn.	<i>Sēmaram karang.</i>
Synacidium horridum L.	<i>Lēpu.</i>
(Unidentified),	<i>L. bērunyut, L. landak.</i>

PLATYCEPHALIDAE (FLAT-HEADS).

Platycephalus punctatus C. V.	<i>Baji-baji.</i>
„ tuberculatus C. V.	„
„ macracanthus Blkr.	„

LEPTOSCOPIDAE (STAR-GAZERS).

Percis pulchella Tem. Schleg.	<i>Pukul gēndang.</i>
--------------------------------------	-----------------------

BATRACHIDAE (FROG-FISHES).

Batrachus grunniens L.	<i>Kērtakok.</i>
-------------------------------	------------------

MASTACEMBELIDAE (SPINY-EELS).

Mastacembelus unicolor C. V.	<i>Tilan or Telan.</i>
„ maculatus C. V.	„ „
„ armatus Lac.	„ „

ANTENNARIIDAE (ANGLER-FISHES).

Antennarius cantori Blkr.	<i>Lēpu, Dēpu, Gēdēmpu.</i>
„ caudimaculatus Günth.	„ „ „
„ coccineus Günth.	„ „ „
„ hispidus El. Schn.	„ „ <i>Kēlalawar.</i>

MALTHIDAE (CROAKERS).

Halieutaea stellata Wahl.	<i>Tēkok.</i>
----------------------------------	---------------

TRIACANTHIDAE (LEATHER-JACKETS).

Triacanthus strigilifer Cant.	<i>Barat-barat.</i>
„ blochii Blkr.	„
„ brevirostris Schleg.	„
„ oxycephalus Blkr.	„ „ <i>Tunjang langit.</i>
„ nieuhoi Blkr.	„
„ spp.	„ <i>Lēmbu.</i>

BALISTIDAE (LEATHER-JACKETS).

Balistes stellatus Lac.	<i>Jëbong, Ayam.</i>
Monacanthus hajam Blkr.	<i>Ayam.</i>
„ sinensis L.	<i>Barat-barat.</i>
„ choerocephalus Blkr.	<i>Kërosok, Ayam.</i>
„ penicilligerus Cuv.	<i>Barat-barat.</i>
„ monoceros L.	<i>Kërosok padi.</i>
Anacanthus barbatus Gray.	<i>Sëligi.</i>

OSTRACIONTIDAE (BOX-FISHES).

Ostracion nasus Bl.	<i>Buntal kotak, B. batu.</i>
„ cubicus L.	„ „ „ „
„ cornutus L.	„ „ „ „

TETRODONTIDAE (GLOBE-FISHES).

Tetrodon lunaris Bl. Schn.	<i>Buntal pisang.</i>
„ reticularis Bl. Schn.	„ <i>duri.</i>

DIODONTIDAE (PORCUPINE-FISHES).

Diodon novemmaculatus Blkr.	<i>Buntal landak.</i>
„ hystrix L.	„ „

SCYLLIIDAE (DOG FISHES).

Scyllium marmoratum Benn.	<i>Yu pendek.</i>
Stegostoma tigrinum L.	<i>Yu chëchak, Yu tokek.</i>
Chiloscyllium indicum L.	<i>Yu tokek, Yu bëlangkas.</i>

CARCHARIIDAE (SHARKS).

Carcharias dussumieri Val.	<i>Yu.</i>
Galeocerdo rayneri McD. B.	<i>Yu lënggiri.</i>
(Unidentified).	<i>Yu jërong, Yu sambaran, Yu punai, Yu laras, Yu chëmangi, Yu bodoh.</i>

SPHYRNIDAE (HAMMER-HEAD SHARKS).

Sphyrna (Zygaena) malleus Risso.	<i>Yu bëngkong, Y. palang, Y. sanggul.</i>
„ (Zygaena) blochii Cuv.	<i>Yu bëngkong, Y. palang, Y. sanggul.</i>

PRISTIDAE (TRUE SAW-FISHES).

Pristis cuspidatus Lath.	<i>Běroi, Yu parang, Yu gěr-gaji, Yu todak.</i>
---------------------------------	---

•• RHINOBATIDAE (BEAKED RAYS).

Rhinobatus thonini Lac.	<i>Yu kia-kia, Kěmějan.</i>
Rhynchobatus djeddensis Forsk.	„ „ „

TORPEDINIDAE (ELECTRIC RAYS).

Narcine timlei Bl. Schn.	<i>Pari kěbas.</i>
Astrape dipterygia Bl. Schn.	„ „ „ <i>P. bunga.</i>
Temera hardwickii Gray.	„ <i>linchin.</i>

TRYGONIDAE (STING-RAYS).

Trygon uarnak Forsk.	<i>Pari kělawar.</i>
„ sephen Forsk.	„ <i>běndera, P. daun.</i>
„ walga M. H.	„ <i>lalat.</i>
„ kuhlii M. H.	„ <i>rimau.</i>
Urogymnus asperrimus Bl. Schn.	„ <i>dědap.</i>

MYLIOBATIDAE (EAGLE-RAYS).

Myliobatis vespertilio Blkr.	<i>Pari kělawar.</i>
„ nieuhofi Bl. Schn.	„ „
Aetobatis narinari Euphr.	„ <i>lang.</i>
Rhinoptera adspersa M. H.	„ <i>daun.</i>
Dicerobatis eregoodoo Cant.	„ <i>paus.</i>
Ceratoptera ehrenbergii M. H.	„ <i>paus, P. kola.</i>

UNIDENTIFIED (RAYS).

Sai, Mengkai, Sětoka.

Authors Consulted.

- BLEEKER, DR. P. Atlas Ichthyologique des Indes Orientales Néer-
landaises. 9 Vols: 1862—1878.
- BOULENGER, DR. G. A. Fishes (Teleostei). }
BRIDGE, DR. T. W. Fishes. }
Cambridge Nat: Hist: Vol. VII. 1904.
- CANTOR, DR. T. Catalogue of Malayan Fishes. Jour: A. S.
Bengal. 1849.
- CLIFFORD, H. AND SWETTENHAM, F. A. Dictionary of the Malay
language. Parts 1-V. 1894—1902.
- DAY, DR. FRANCIS. Fauna of British India. Fishes. 2 Vols:
1889.
- DUNCKER, GEORG. Die Fische der Malayischen Halbinsel. 1904.
- GUNTHER, DR. ALBERT. Study of Fishes. 1880.
- MEEK, DR. The Migrations of Fish. 1916.
- ROUGHLEY, T. C. Fishes of Australia and their Technology.
1916.
- SEALE, ALVIN. Fishery resources of the Phillipines. (*Philippine
Jour: Science*. 1908).
- STEAD, DAVID A. Fishes of Australia. 1906.
- STEAD, DAVID A. Edible Fishes of New South Wales. 1908.
- STEAD, DAVID A. A brief review of the Fisheries of New South
Wales. 1910.
- TENISON-WOODS, REVD. J. E. Fish and Fisheries of New South
Wales. 1882.
- WEBER AND DE BEAUFORT, DRS. The Fishes of the Indo-Austra-
lian Archipelago. 3 Vols: 1911—1913.
- WILKINSON, R. J. Malay-English Dictionary. 2 Vols: 1901.

Index to English Names

A.

Anchovy 18, 60.
Angler-Fishes 49, 99.

B.

Barracouta 71, 76, 84.
Barracuda 27, 91.
Barramundi 32.
Bar-tailed Goat-Fish 36.
Batfish, Dark 38.
Batfish, Silvery 37, 62.
Bat-Fishes 37, 95.
Beaked-Ray 52, 101.
Big-eyed Herring 15.
Black-spotted Rock-Cod	.. 32.
Black Trevally 38, 96.
Blow-pipe Fishes 31, 91.
Blue-spotted Groper 42.
Bombay Duck 19, 72.
Bony Bream 18.
Box-Fishes 50, 100.
Bream, Sea 35, 94.
Bream, Silver 34, 93.
Brown-spotted Hind 32.
Bull's-Eyes 30, 91.
Butter-Fish 37, 95.

C.

Carp 10, 20, 88.
Catla 22.
Cat-fish 19, 87.
Climbing Perch 40, 60, 77.
Conger-Eels 22, 89.
Coral-Fishes 37, 41, 95, 96
Croakers 49, 99.

D.

Dart 43, 74.
Devil-fishes 53.
Diamond-scaled Mullet	.. 26.
Dog-Fishes 50, 100.
Dorab 74.
Drummers 30, 91.
Dusky-Perch 30, 91.

E.

Eagle-Rays 53, 101.
Eels 7, 22.
Eels, Conger 22, 89.
Eels, Spiny 49, 99.
Eels, Swamp 22, 89.
Electric Rays 52, 101.

F.

Feather-Backs 15, 86.
Fighting-Fishes 40, 96.
Flat-Fishes 45, 98.
Flat-heads 48, 99.
Flounder 46.
Flying-Fishes 23, 90.
Fresh-water Perches 31, 92.
Frog-Fishes 49, 99.

G.

Garfishes 23, 90.
Gar-Pike 23, 90.
Giant-Herrings 15, 86.
Globe-Fishes 50, 100.
Goat-Fish, Bar-tailed 36.
Gobies 46, 98.
Goblin-Fishes 48, 99.
Gourami 38, 96.
Grey-Mullets 24, 90.
Groper, Queensland 32.
Grunters 35, 94.

H.

Hair-back 18.
Hair-tails 45, 98.
Half-beak 23.
Halibut, Queensland 46.
Hammer-head Sharks 50, 100.
Herrings 17, 86.
Hilsa 17.
Horse-Mackerels 42, 97.

J.

Jew-Fishes 34, 93.
------------------	------------

K.

Knight-Fish 30, 91.
-------------------	------------

L.

Leather-jackets 49, 99, 100.
Loaches 20, 88.

M.

Macassar Redfish 17.
Mackerel 44, 97.
Mackerel, Horse 42, 97.
Mackerel, Spanish 44.
Mahseer 21.
Milk-Fish 15, 86.
Millions 19, 87.

Moon-Fish 38, 62, 95.				S.				
Moon-Fish, Silvery 43, 61.				Sable-Fish 17.			
Mud-skipper 47, 82.				Sail-Fishes 45, 98.			
Mullet, Diamond-scaled 26.				Salmon-Herring 15, 56.			
Mullet, Grey 24, 90.				Sand-Smelts 23, 90.			
Mullet, Red 36, 95.				Saw-Fishes 51, 101.			
Murrel 28, 91.				Sea-Horses 22, 90.			
					Sea-Perches 31, 92.			
N.					Selemah 34, 78, 94.			
Nannygai 29, 91.				Shad 17, 86.			
					Sharks 50, 100.			
O.					Sharks, Hammer-head 50, 100.			
Ox-Eye 15				Silver-Bream 34, 93.			
					Silvery Moon-Fish 43, 61.			
P.					Skeleton-Fishes 22, 90.			
Paradise-Fish 41.				Smelt, Queensland 19, 87.			
Parrot-Fishes 41, 96.				Smelt, Sand 23, 90.			
Parrot-Wrasses 42, 97.				Snappers 33, 92.			
Perch, Climbing 40, 60, 77.				Spanish Mackerel, Barred 44.			
Perches, Freshwater 31, 92.				Spiny-Eels 49, 83, 99.			
Perches, Sea 31, 92.				Star-Gazers 49, 76, 99.			
Pescados del Rey 24.				Sucking-Fishes 48, 98.			
Pike, Short-finned 27.				Swamp-Eels 22, 89.			
Pomfrets 28, 91.								
Porcupine-Fishes 50, 100.				T.				
					Tarpon 15.			
Q.					Tarwhine 36.			
Queen-Fish 43.				Ten-pounder 15.			
Queensland Groper 32.				Thread fins 27, 91.			
Queensland Halibut 46.				Trevally, Black 38, 96.			
Queensland Smelt 19, 87.				Trumpeter, Queensland 35.			
Queensland Trumpeter 35.				Trumpeter Whiting 33.			
					Tunny 44, 97.			
R.					W.				
Rays, Beaked 52, 101.				Whitebait 17.			
Rays, Eagle 53, 101.				White-Mullet 15.			
Rays, Electric 52, 101.				Whiting 33, 93.			
Rays, Sting 52, 101.				Wrasses 41, 96.			
Red-Mullet 36, 95.								

ADDENDA ET CORRIGENDA.

P. 7: for Tarpin read Tarpon.

P. 59: add **Bēras-bēras**. *Sparus sarba*.

The Tarwhine of Australia.

Sea-Bream. Fam. SPARIDAE.

P. 95: add **Sparus sarba** Forsk. *Bēras-bēras*.

LIST OF PLATES.

Plate.		
I.	BULAN	Ox-Eye, or Big-eyed Her- ring.
II.	BĒLIDA	Feather-back.
III.	PARANG-PARANG	The Dorab.
IV.	SĒLANGAT	Hairback or Gizzard Shad.
V.	TĒRUBOK KORIN	The Hilsa.
VI.	TAMBAN PANJANG	"Sardine."
VII.	TAPAH	River Cat-fish.
VIII.	BĒGAHAK or LAIS	River Cat-fish.
IX.	JAHAN	Sea Cat-fish.
X.	KĒLAH	Carp.
XI.	TĒMBĒLIAN	Carp.
XII.	UMBU'T-UMBU'T	Carp.
XIII.	HWAN or CHOW	Chinese Carp.
XIV.	LIAN or LIN	Chinese Carp.
XV.	MALONG	Conger-eel.
XVI.	TODAK	Gar-Pike.
XVII.	PUPUT or JOLONG- JOLONG	Gar-fish.
XVIII.	BĒLANAK ANDING	Grey-Mullet.
XIX.	BĒLANAK TAMOK	Diamond-scaled MuHet.
XX.	KURAU	Mango fish.
XXI.	ALU-ALU	Barracuda.
XXII.	BAWAL PUTEH	Pomfret.
XXIII.	ARUAN	Murrel.
XXIV.	SĒBĒKAH KARANG	"Nannygai."
XXV.	SUMPIT-SUMPIT	Blow-pipe fish.
XXVI.	KERONG-KERONG	Sea-Perch.
XXVII.	KĒRAPU	Sea-Perch.
XXVIII.	SIAKAP	Sea-Perch.
XXIX.	IKAN MERAH	Snapper.
XXX.	BULUS-BULUS	"Whiting."
XXXI.	TĒMBĒREH	Jew-fish.
XXXII.	GĒLAMA TIKUS	Jew-fish.
XXXIII.	KAPAS-KAPAS	"Silver-Bream."
XXXIV.	GĒRUT-GĒRUT	Grunter.
XXXV.	PASIR-PASIR	Sea-Bream.

XXXVI.	DĚLAH	Sea-Bream.
XXXVII.	ASOH-ASOH	Sea-Bream.
XXXVIII.	PINANG-PINANG	Sea-Bream.
XXXIX.	BIJI NANGKA	Red mullet.
XL.	GĚDABANG	Silver Bat-fish.
XLI.	KETANG	Spotted Butter-fish.
XLII.	BABI	Coral-fish.
XLIII.	TUDONG PĚRIOK	Coral-fish.
XLIV.	DAUN BAHARU	Moon-fish.
XLV.	DĚNGKIS	"Black Trevally."
XLVI.	DEBAM	"Black Trevally."
XLVII.	KALUI	Gurami.
XLVIII.	GOMBING	Coral-fish.
XLIX.	TOKAK	Wrasse.
L.	BĚCHOK	Parrot-Wrasse.
LI.	CHĚNCHARU	Horse-Mackerel (Hard tail)
LII.	DAING BĚLANG	Banded Horse-Mackerel.
LIII.	CHERMIN	Silvery Moon-fish.
LIV.	SAGAI	Trevally.
LV.	NYIOR-NYIOR	Dart.
LVI.	TALANG	Queen-Fish.
LVII.	TONGKOL	Tunny.
LVIII.	TONGKOL CHORENG	Tunny.
LIX.	TĚNGGIRI PAPAN	Spotted Spanish Mackerel.
LX.	TĚNGGIRI BATANG	Barred Spanish Mackerel.
LXI.	SĚLAYUR	Hair tail.
LXII.	SA-BĚLAH	"Halibut."
LXIII.	LIDAH	"Sole."
LXIV.	BAJI-BAJI	Flat-head.
LXV.	BARAT-BARAT	Leather-jacket.
LXVI.	JĚBONG	Leather-jacket.
LXVII.	YU TOKEK	Dog-fish.
LXVIII.	YU PALANG	Hammer-head Shark.
LXIX.	YU KĚMĚJAN	Beaked-Ray.
LXX.	PARI BĚTING	Sting-Ray.
LXXI.	KĚRAPU, KURAU, TULANG, PARANG	MERAH, TĚNGGIRI,
LXXII.	YU, PARI, MALONG, DURI.	.

PLATE I



0 1 2 3 4 5 6 inches

BULAN

Ox - Eye. or Big eyed Herring.

(*Megalops cyprinoides*).

PLATE II



BĚLIDA

Feather - back.

(*Notopterus notopterus*).

PLATE IV



SĚLANGAT

Hairback or Gizzard Shad

(*Dorosoma nasus*).

PLATE V




0 1 2 3 4 5 6 7 8 9 10 11 12 inches

TĚRUBOK KORIN

The Hilsa

(*Clupea (Alosa) sp.*)

PLATE VI



0 1 2 3 inches

TAMBAN PANJANG

"Sardine"

(*Clupea* sp.)

PLATE VII



0 1 2 3 4 5 6 7 8 9 10 11 12 INCHES

TAPAH

River Cat-fish.

(*Wallago sp.*)

PLATE VIII



0 1 2 3 4 5 6 7 8 9 10 11 12 INCHES

BĚGAHAK or LAIS

River Cat-fish.

(*Belodontichthys dinema*)

PLATE IX



0 1 2 3 4 5 6 7 8 9 10 11 12 INCHES

JAHAN

Sea Cat-fish.

(*Arius* sp.)

PLATE X



0 1 2 3 4 5 6 7 8 9 10 11 12 INCHES

KĚLAH

Carp.

(*Barbus sp.*)

PLATE XI



TĚMBĚLIAN

Carp.

(*Barbus* sp.)

PLATE XII



UMBUT-UMBUT

(*Barynotus microlepis*)

Carp.

PLATE XIII




HWAN or CHOW

Chinese Carp.

(*Otenopharyngodon idellus*)

PLATE XIV



0 1 2 3 4 5 6 7 8 9 10 11 12 inches

LIAN or LIN

Chinese Carp.

(*Thymichthys* sp.).

PLATE XV




MALONG

Conger-eel.

(*Muraenesox* sp.)

PLATE XVI



0 1 2 3 4 5 6 7 8 9 10 11 12 INCHES

TODAK

Gar Pike

(*Belone sp.*).

PLATE XVII



PUPUT or JOLONG-JOLONG

Gar-fish.

(*Hemirhamphus far*).

PLATE XVIII



BĚLANAK ANDING

Grey Mullet.

(*Mugil borneensis*).

PLATE XIX



0 1 2 3 4 5 6 Inches

BĒLANAK TAMOK

Diamond-Scaled Mullet.


(*Mugil waigiensis*).



BĒLANAK TAMOK

Diamond-Scaled Mullet.

(*Mugil waigiensis*).





ALU-ALU

Barracuda.

(*Sphyræna commersonii*.)

PLATE XXII



BAWAL PUTEH

Pomfret.


(*Stromateus cinereus*).



ARUAN

Murrel.

(*Ophiocephalus* sp.).



0 1 2 3 Inches

SĚBĚKAH KARANG

"Nannygai".


(*Mogripristis murdjan*).



SUMPIT-SUMPIT

Blow-pipe fish.


(*Toxotes jaculator*).



KERONG-KERONG

Sea-Perch.

(*Therapon* sp.). ?



KĚRAPU

Sea-Perch.

(*Epinephelus* sp.).



SIAKAP

Sea-Perch.

(*Lates calcarifer*).



IKAN MERAH

Snapper.

(*Lutianus* sp.).

PLATE XXX



BULUS-BULUS

"Whiting".

(*Sillago sihama*).



TĚMBĚŘEH

Jew-fish.

(*Sciaena* sp.)

PLATE XXXII



GĒLAMA TIKUS

Jew-fish.

(*Sciaenidae* sp.).



KAPAS-KAPAS

"Silver-Bream".

(*Gerres abbreviatus*).



GĚRUT-GĚRUT

Grunter.

(*Pristipoma hasta*).




PASIR-PASIR

Sea-Bream.

(*Scolopsis sp.*). ;

PLATE XXXVI



DÉLAH

Sea. Bream.

(*Caesio kneri*).




ASOH-ASOH

Sea Bream.

(*Lethrinus nebulosus*).

PLATE XXXVIII



0 1 2 3 4 5 6 inches

PINANG-PINANG

Sea-Bream.

(*Sparus datnia*).

PLATE XXXIX



BIJI NANGKA

Red mullet.

(*Upeneus tragula*).



GĔDABANG

Silver Bat-fish.

(*Pseittus argenteus*).

PLATE XLI



KETANG

Spotted Butter-fish.

(*Ephippus argus*).

PLATE XLII



BABI

Coral-fish.

(*Holacanthus* sp.).

PLATE XLIV



0 1 2 3 4 5 6 7 8 9 10 11 12 Inches

DAUN BĀHARU

Moon-fish.

(*Drepanè punctata*).



PLATE XLVI




DEBAM

Black Trevally.

(*Teuthis java*).

PLATE XLVII




KALUI

Gurami.

(*Osphromenus olfax*).

PLATE XLVIII



GOMBING

Coral-fish.

(*Glyphidodon* sp.)

PLATE XLIX



TOKAK

Wrasse.

(*Choerops* sp.).

PLATE L



BĚCHOK

Parrot-Wrasse.

(*Pseudoscarus* sp.).




0 1 2 3 4 5 6 inches

CHĒNCHARU

Horse-Mackerel (Hard tail)

(*Caranx rottileri*).



DAING BÉLANG

Banded Horse-Mackerel.

(*Caranx speciosus*).

PLATE LIII




CHERMIN

Silvery Moon-fish.

(*Caranx gallus*).

PLATE LIV



SAGAI

Trevally.

(*Caranx* sp.).




PLATE LVI



0 1 2 3 4 5 6 7 8 9 10 11 12 Inches

TALANG

Queen-fish.

(*Chorinemus moadetta*).



TONGKOL

Tunny.

(*Thynnus* sp.).



0 1 2 3 4 5 6 7 8 9 10 11 12 inches

TONGKOL CHORENG

Tunny.

(*Thynnus thunnus*).

PLATE LIX



0 1 2 3 4 5 6 7 8 9 10 11 12 INCHES

TENGGIRI PAPAN

Spotted Spanish Mackerel.

(*Cybium guttatum*).

PLATE LX



TENGGIRI BATANG

Barred Spanish Mackerel.

(*Cybium commersonii*).

PLATE LXI



0 1 2 3 4 5 6 7 8 9 10 11 12 INCHES

SĚLAYUR

Hair-tail.

(*Trichurus haumela*).

PLATE LXII



SA-BĚLAH

"Halibut".

(*Psettodes crumei*).

PLATE LXIII



0 1 2 3 4 5 6 Inches

LIDAH

"Sole"

(*Synaptura sp.*).

PLATE LXIV



BAJI-BAJI

Flat-head.


(*Platycephalus* sp.).



BARAT-BARAT

Leather jacket.

(*Monacanthus* sp.).



0 1 2 3 4 5 6 inches



YU TOKEK

Dog-fish.

(*Scyllium* sp.).

PLATE LXVIII



YU PALANG

Hammer-head Shark.

(*Sphyrna (zygaena) sp.*).



YU KĚMĚJAN

Beaked-Ray.

(*Rhynchobatus djeddensis*).

PLATE LXX



PARI BĚTING

Sting-Ray.

(*Trygon uarnak*).


PLATE LXXI



KĒRAPU, KURAU, -MERAH, TĒGGIRI,
TALANG, PARANG.

Photo taken at Clyde Terrace Market Singapore

PLATE LXXII



YU, PARI, MALONG, DURI.

Photo taken at Clyde Terrace Market Singapore

